

# AN INFORMATION–THEORETIC APPROACH TO PARTIALLY IDENTIFIED AUCTION MODELS\*

SUNG JAE JUN<sup>¶</sup> AND JORIS PINKSE<sup>§</sup>

Department of Economics  
The Pennsylvania State University

March 24, 2021<sup>§</sup>

We consider a situation in which we have data from ascending auctions with symmetric bidders, independent private values, and exogenous entry in which the bidders' value distribution is partially identified. Focusing on the case in which the seller intends to use a second price auction, we discuss how to determine an optimal reserve price. We justify the use of maximum entropy, explore the properties of the estimand, determine the asymptotic properties of our maximum entropy estimator, evaluate its behavior in a simulation study, and demonstrate its use in a modest application. As an extension, we propose a maxmin decision rule with entropy regularization, which includes Aryal and Kim (2013) and the maximum entropy solution as extreme cases.

**Keywords:** English auctions, optimal mechanism, partial identification, maximum entropy, inequality constraints, and nonparametric inference.

**JEL:** C01; C10; C57.

---

\*We thank Andres Aradillas–Lopez, Gaurab Aryal, Michael Gechter, Ed Green, Nima Haghpanah, Phil Haile, Marc Henry, Kei Hirano, Dong–Hyuk Kim, Vijay Krishna, Laurent Lamy, Brad Larsen, Peter Newberry, Francesca Molinari, Ran Shorrer, Guofu Tan, the coeditor, the associate editor, three referees, and participants of the SVEC conference in Vancouver, the conference in memory of Art Shneyerov in Paris, the KER International conference in Seoul, the PSU–Cornell econometrics and IO conference, the EARIE meeting in Barcelona, and departmental seminars at Penn State and the University of British Columbia for their helpful comments.

<sup>¶</sup>619 Kern Bldg, University Park, PA 16802, sjun@psu.edu

<sup>§</sup>joris@psu.edu

<sup>§</sup>The first circulated version was on May, 2017.

# 1. Idea and context

We consider English auctions with exogenous entry and symmetric bidders under the independent private values (IPV) paradigm in which the number of potential bidders  $n$  and some bids are observed. Since this is an English auction, there is no one-to-one mapping between bids made by a given bidder and his valuation as there would be in a symmetric sealed bid IPV auction without a reserve price. Indeed, bidders can continue to submit bids after they have submitted one or may not submit a bid at all because the current bid level already exceeds their private value.

We assume that bidders can only select bids from a prespecified finite set. We focus on the simplest case in which bids belong to  $\{\Delta, 2\Delta, \dots\}$  with  $\Delta > 0$ , but this is not essential. Haile and Tamer (2003, HT) consider the possibility that there is a minimum positive bid increment, but HT's results apply in our environment, also. Like HT, we allow for jump bidding. From here on, we will discuss HT's methodology in the context of our environment. If  $\Delta > 0$  then the bidders' value distribution is only partially identified.<sup>1</sup> Indeed, HT obtain bounds on the unknown distribution function  $\mathcal{F}$  of private values which, both in HT and here, is assumed to be continuous.<sup>2</sup> HT then add a strict pseudoconcavity assumption on the *pseudoprofit* function  $\mathcal{Q}(r) = r\{1 - \mathcal{F}(r)\}$  to obtain bounds on the optimal reserve price.<sup>3</sup>

Like HT, we think about the problem of setting an optimal reserve price in a hypothetical second price auction with  $\Delta = 0$ . The rationale for this is, as HT have shown, that an English auction with a reserve price and no minimum bid increment is revenue-equivalent to a second price sealed bid auction with the same reserve price under the behavioral assumptions made in HT, as long as the English auction can be represented by a “feasible auction mechanism.”<sup>4</sup> However, the revenue equivalence theorem (Myerson, 1981) does *not* imply that a second price auction is optimal. We revisit these issues in section 2 and appendix A.

Although there is precedence for HT's pseudoconcavity assumption (which we do not need) in the literature, there are many plausible value distributions that do not satisfy it and there is no clear

---

<sup>1</sup>Athey and Haile (2002) contains identification results on English auctions.

<sup>2</sup>In view of Chesher and Rosen (2018), HT's bounds are not entirely sharp because they do not exploit correlation among bids. This is however irrelevant for the main ideas put forth in this paper. We thank Laurent Lamy and Francesca Molinari for pointing this out to us.

<sup>3</sup>The pseudoprofit function is the seller's profit as a function of the reserve price if there is only one bidder. We set the seller's value to zero for presentational simplicity.

<sup>4</sup>A feasible auction mechanism requires a Nash equilibrium in the bidding game (Myerson, 1981, p62). The case  $\Delta = 0$ , both here and in HT, rules out the bulk of situations giving rise to jump bidding in equilibrium. For instance, Avery (1998) requires affiliation, whereas we follow HT in assuming independence. Further, Daniel and Hirshleifer (1998) is mostly about a scenario in which bidding is costly. Daniel and Hirshleifer (1998) do provide an example of a Nash equilibrium with zero costs, no minimum bid increment, and symmetric bidders in which there exists an equilibrium with jump bidding, but it is both fragile and does not satisfy HT's requirement that bidders will not let rivals win at a price that they are willing to beat. In fact, we are not aware of an example in which there are jump bids in equilibrium with symmetric bidders, exogenous entry, independent private values, no minimum bid increment, and in which HT's behavioral assumptions are satisfied.

path to usable bounds on the optimal reserve price in a second price auction from bounds on  $\mathcal{F}$  without it: example 1 in section 2 and figures 5 and 6 in section 4 contain such distributions.

Further, bounds on the optimal reserve price are interesting, but they are of limited use to a seller faced with the problem of choosing a reserve price: she will still have to pick a single number. Indeed, as we show in section 4, if the number of bidders is small then expected revenue can vary substantially with the choice of reserve price, even if one were only to consider reserve prices that belong to the HT identified set. The question of interest in this paper, then, is how to pick a reserve price if the value distribution  $\mathcal{F}$  is only partially identified.

This question already has an answer in Aryal and Kim (2013, AK) in the somewhat different case in which only the winning bid is observed. Their idea can be summarized as follows. AK assume that the winning bid is no greater than the highest value. So, the  $n$ -th power of  $\mathcal{F}$  is bounded above by the distribution function of the winning bid. They further assume that the seller is ambiguity-averse (Gilboa and Schmeidler, 1989), but *not* risk-averse, which means that the seller's objective is to maximize minimum expected revenue over all value distributions.<sup>5</sup> AK's results show that if candidate distribution  $\mathcal{F}_1$  stochastically dominates alternative  $\mathcal{F}_2$  then the expected revenue associated with  $\mathcal{F}_1$  is no less than that with  $\mathcal{F}_2$  for any reserve price. Therefore, a lower bound to expected revenue obtains when the value distribution is set equal to its upper bound. In other words, the value distribution  $\mathcal{F}_{AK}$  that corresponds to the smallest expected revenue is such that  $\mathcal{F}_{AK}^n$  equals the distribution of the winning bid. Therefore, the seller can choose a reserve price as though  $\mathcal{F}_{AK}$  were the true value distribution.

The methodology we propose can be used in the environment analyzed in AK, but we choose to develop our methodology in an environment similar to HT's world which is methodologically more interesting for our approach: our methodology uses both the upper and the lower bounds of the identified set but in AK's environment the lower bound is always trivial.<sup>6</sup> AK's methodology can be extended to HT's case, also. Indeed, one can simply look at the worst possible value distribution that is consistent with HT's bounds, i.e. the HT upper bound of the value distribution. As far as we know, this has not been done, but it is not difficult. From here on, we will take that leap and discuss AK's approach in the context of HT's conditions only.

A more general way of addressing the reserve price selection problem is to express it in terms of a decision-theoretic framework by assuming that a seller chooses the reserve price to maximize minimum expected revenue where the minimum is taken over all value functions  $\mathcal{F}$  that belong to the HT identified set and whose (relative) entropy is no less than some lower bound  $\mathcal{E}^*$  specified by the seller. In other words, we cast the seller's decision problem in the same framework as AK but we impose an extra entropy regularization condition; without the regularization AK's solution

---

<sup>5</sup>She is not risk-averse, because she is still trying to maximize expected revenue instead of the expectation of a strictly concave function of revenue. In AK, the seller is ambiguity-averse because she does not like the ambiguity resulting from the fact that the value distribution is unknown.

<sup>6</sup>Further, the upper bound of the identified set in AK's environment is greater than that in HT's.

exclusively depends on the upper bound of the distribution function of valuations. With this setup, the choice of  $\mathcal{E}^*$  reflects the seller's attitude about ambiguity and plausibility: she balances an aversion to ambiguity with a desire to rule out distributions she considers implausible as measured by the entropy. It allows her to provide a guess of the value distribution known as a *reference distribution* (which we take to be a uniform) and only permit deviations within a certain distance from the reference distribution, where the distance is measured by (minus) the (relative) entropy, i.e. the Kullback–Leibler divergence criterion. There is hence a *shrinkage* aspect to this approach in that (absent point identification in the original problem) the choice of distribution function is shrunk towards the reference distribution. The value of  $\mathcal{E}^*$  determines the tradeoff between ambiguity aversion and plausibility: for  $\mathcal{E}^* = -\infty$  one obtains AK's pure maxmin approach and with  $\mathcal{E}^*$  chosen maximally one obtains *maximum entropy*, a method from the *information theory* literature (see e.g. Topsøe (1979), Harremoës and Topsøe (2001), Grünwald and Dawid (2004, GD)); the fact that the solution is sensitive to the choice of  $\mathcal{E}^*$  is a natural consequence of partial identification. A detailed discussion of this entropy–regularized decision–theoretic approach can be found in appendix B. In the main text we consider the special case in which the seller chooses  $\mathcal{E}^*$  maximally, i.e. uses the maximum entropy principle. Below we discuss the choice of a reference distribution and explain why the (maximum entropy) special case is particularly interesting.

We start with the choice of a reference distribution. As noted, one possibility is to ask the seller for a guess of what the true value distribution looks like. The need to pick a reference distribution can be compared to the need in Bayesian decision theory to specify a prior distribution, but the demand here is less onerous: Bayesian decision theory would require the seller to specify a prior over the set of value distributions, which is probably an unreasonable amount of information to elicit from a seller.

A second possibility is to start with a more restricted model, in which there is point identification, e.g. by adding behavioral assumptions. One can then estimate the restricted model which produces an estimated value distribution which can be used as a reference distribution in our procedure, albeit that the inference methods proposed in this paper would need to be extended to accommodate sampling errors in the reference distribution.

There are two reasons for setting the reference distribution to a uniform as the default. First, it simplifies notation noting that the extension to other reference distributions is tedious but straightforward, albeit that the maximum entropy solution will then not be piecewise linear, as we will show it is here.<sup>7</sup> Second, it is a natural choice in the information theory framework since it means that we would be maximizing Shannon's entropy subject to the identified bounds of the value distribution. Note that the uniform distribution reflects our *ex ante* 'ignorance' about what the value distribution should look like, which is in line with the rationale of *Occam's razor*.

---

<sup>7</sup>The maximum entropy distribution function will be piecewise linear; the corresponding density function hence piecewise constant.

We now discuss why the maximum entropy solution is of particular interest. We offer three arguments. First, the maximum entropy principle can be thought of as a special case of robust control theory (e.g. Hansen and Sargent, 2008) and it can be understood as a minmax decision with a specific loss function. Indeed, Topsøe (1979), Harremoës and Topsøe (2001), and GD *inter alia* provide a justification of the maximum entropy principle from the usual minmax perspective of decision theory. GD is particularly relevant because they show that choosing an entropy-maximizing distribution from a set  $\mathcal{C}$  is equivalent to selecting a robust Bayes act against  $\mathcal{C}$  (or  $\mathcal{C}$ -minmax act) for the log loss decision problem. Indeed, in a statistical game in which a decision maker specifies a distribution from  $\mathcal{C}$  and nature reveals values from an unknown distribution (that may or may not belong to  $\mathcal{C}$ ), maximizing entropy over  $\mathcal{C}$  is a dual decision to considering the worst-case expected log loss among  $\mathcal{C}$  and minimizing it. In our context,  $\mathcal{C}$  corresponds to the identified set of the value distributions, albeit that the GD results do not cover the case of inequality restrictions.<sup>8</sup> One could argue that this statistical game is not directly relevant to the seller, who only cares about expected profits. But as argued earlier, the maximum entropy solution can be justified from the perspective of expected profit maximization, also, by introducing entropy-based regularization; see appendix B. This approach is similar in spirit to Hansen and Sargent (2008), who consider a constrained minmax approach, where entropy is used to control for the admissible level of misspecification.

Second, it is well-known in the information theory literature that the maximum entropy principle constitutes the unique rule of inference that satisfies a certain set of axioms (e.g. Golan, 2018). The axioms are as follows: (i) the solution must be unique; (ii) the solution must be invariant to the transformation of the coordinate system; (iii) breaking the problem into smaller pieces should not change the solution; i.e. solving two problems on  $[0, 1/2]$  and  $[1/2, 1]$  separately should yield the same solution as solving one problem on  $[0, 1]$ ; (iv) solving for a joint distribution or for a marginal and a conditional distribution should yield the same answer; (v) without extra information, the solution should be the same as the prior. So, in the present context, the maximum entropy value distribution  $\mathcal{F}^*$  is the only solution that satisfies all five axioms.

Third, the maximum entropy principle has several practical advantages. For instance, it is easy to implement, offering a solution that is analytically tractable. Further, it is straightforward to add any additional information or insights the seller might have, or indeed information or insights that can be inferred from the seller's actions, such as the reserve price she chose to set in the observed auctions.<sup>9</sup>

Our maximum entropy method yields a unique solution  $\mathcal{F}^*$  that satisfies the HT distribution

---

<sup>8</sup>It is outside the scope of this paper to generalize the GD theory to the case of inequality restrictions.

<sup>9</sup>For instance, if  $\Delta = 0$  and  $\mathcal{Q}$  is strictly pseudoconcave then  $r^*$  satisfying  $\mathcal{Q}'(r^*) = 0$  would be the unique optimal reserve price. So, when the reserve price  $r^*$  the seller chose to set is observed, imposing  $\mathcal{Q}'(r^*) = 0$  in the maximum entropy optimization problem could be a reasonable, albeit imperfect, choice. We thank Peter Newberry for this suggestion.

function bounds and is piecewise linear.<sup>10</sup> The maximum entropy value distribution  $\mathcal{F}^*$  then implies an optimal reserve price  $r^*$ , which is necessarily between the HT bounds whenever HT’s strict pseudoconcavity assumption is satisfied. However, the maximum entropy distribution itself need *not* satisfy the pseudoconcavity assumption.

Neither AK’s methodology nor our maximum entropy approach requires the pseudoconcavity assumption to select a reserve price. However, absent strict pseudoconcavity, Myerson (1981)’s regular case assumptions are violated and a second price auction and Myerson’s optimal auction may no longer be revenue–equivalent.

Having the maximum entropy value distribution in hand, one can use it to implement Myerson’s optimal auction mechanism instead of simply choosing a reserve price in a second price auction,<sup>11</sup> which is not possible with either HT or AK. Indeed, we discuss optimal auctions in appendix A. However, as noted in appendix A, the gain from doing this is typically small and Myerson’s mechanism is likely to be too cumbersome to implement in a real world environment. Hence, we focus on second price auctions in the body of the paper.




Figure 1: Comparison of AK’s and ME’s choice of the value distribution when  $n = 2$ ,  $\Delta = 0.1$ , and the true value distribution is uniform. The left panel shows the distribution functions and the right panel shows the probability mass function of AK and the probability density function of ME.

Both AK and our paper use HT’s framework to obtain bounds on the value distribution, but it is largely irrelevant for our approach how the bounds are obtained. Indeed, the ME principle we advocate can be applied whenever the distribution function of interest is only partially identified. For example, Coey et al. (2017) have shown how to bound the value distribution function when

<sup>10</sup>If the seller had additional information that was added as restrictions then the ME solution might not be piecewise linear.

<sup>11</sup>Myerson assumes continuity of the value density function, but his results extend to our discontinuous maximum entropy value density.

bidders do not have to be symmetric. This is a different framework from HT's, and hence AK's monotonicity results do not apply. Further, auctions and reserve prices are not the only examples where partial identification of a distribution function arises. For instance, Larsen (2014) studies bargaining games in the used car industry. Larsen obtains partial identification of the seller's value distribution. He then considers welfare measures to investigate the bargaining efficiency problem. In principle, one can consider maximizing the minimum welfare, but some of the measures lack the monotonicity property that makes AK's solution computationally feasible. In contrast, the ME principle remains simple to apply: it can serve as a useful summary in the discussion of optimal market designs.

Aside from the philosophical differences between AK's approach and ours, it is evident that AK's approach leads to simpler estimation and inference procedures since one only needs to estimate a distribution of observables (the winning bids) nonparametrically. On the other hand, the reason that our approach leads to more complicated estimation and inference procedures is, in part, that we use more information. Indeed, HT use multiple distribution function bounds and pick the tightest ones.<sup>12</sup> With AK's approach (in our environment) one would, as noted, only use the HT upper bound on the distribution function. Maximum entropy uses the HT upper and lower bounds, plus an information criterion. Indeed, maximum entropy entails a constrained optimization problem, which leads to interesting inference problems as will become apparent below. Further, the entropy-maximizing value distribution is continuous, whereas the value distribution used for AK's reserve price is discrete, or perhaps more precisely: the AK value distribution results from the discrete limit of a sequence of continuous value distributions: the value distribution must be continuous for the bounds in HT to apply. Figure 1 shows an example, in which the true value distribution is uniform.

Another difference between AK's approach and ours arises when the reserve price in the observed auctions is binding: the AK approach *never* results in an optimal reserve price that is less than what was used in the data used for estimation; this is in contrast to our approach.<sup>13</sup> If there are many auctions without positive bids in the data then AK's optimal reserve price is the same as the reserve price observed in the data,<sup>14</sup> which we think is extreme. In contrast, the maximum entropy optimal reserve price can be as small as half the reserve price used in the data. We discuss this issue in more detail in section 2.

But neither method generally dominates the other or, put differently, either method is better than the other depending on which property is deemed preferable. If the seller is ambiguity-averse but not risk-averse then AK is the method of choice. In other cases, the maximum entropy has a lot going for it. Either method can produce a higher expected revenue under different circumstances. Indeed, these are not the only two possibilities. In other work (Jun and Pinkse, 2016) and a different problem,

---

<sup>12</sup>These are essentially 'intersection bounds.'

<sup>13</sup>We thank Andres Aradillas-Lopez and Vijay Krishna for suggesting that we look into the effect of the reserve price used in the data on results.

<sup>14</sup>We are using a single reserve price here, but similar issues arise when the reserve price in the data varies by auction.

we pursue the possibility of imposing a (Dirichlet process) prior on the class of distribution functions, but that approach comes with unpleasanties: it is painful to obtain a solution and it requires the arbitrary choice of input parameters. Since different reserve prices are optimal for different value distributions (consistent with the data), which method is better is a matter of philosophy, not econometrics. In section 4 we explore some differences between AK’s approach and ours in some examples.

In these examples we simulate the *population* and consider both ‘regular’ and ‘irregular’ cases, i.e. cases in which HT’s strict pseudoconcavity assumption is satisfied and cases in which it is violated. In the regular cases our results are not surprising: HT’s bounds for the optimal reserve price were well-defined but they are too wide to help the seller select a reserve price. Both AK’s solution and ours are always contained in the HT bounds in the regular case. For the irregular case, we consider two different scenarios: one where the value distribution is regular and unimodal but the function  $\mathcal{Q}$  has a flat area, and the other in which the value distribution is bimodal and  $\mathcal{Q}$  has two distinct local maxima.

HT’s results do not apply in the irregular case, but if one nevertheless follows their procedure here one obtains uninformative bounds or nonconvex identified sets for the optimal reserve price, depending on the shape of the  $\mathcal{Q}$  function. Both AK’s approach and ours produce a point decision, but neither dominates the other in terms of true expected revenue, which is in line with the theory. Again, the choice is between insurance against the worst case expected revenue (AK) and the most ‘reasonable’ value distribution as defined by the entropy criterion (JP).

The discussion has thus far focused on the case in which the distributions of observables are known, in which case  $\mathcal{F}^*$  and  $r^*$  can be determined. In practice, however,  $\mathcal{F}^*$ ,  $r^*$ , and the corresponding expected revenue would have to be estimated. Addressing issues resulting from estimation error can be relevant to the seller, also. She may, for instance, only wish to deviate from established practice if the ME reserve price is significantly different from the one she currently uses, or indeed if the corresponding expected revenue is significantly higher. For the purpose of inference, we assume that the HT bounds can be  $\sqrt{N}$ -consistently estimated, and we express the distribution of the objects of interest in terms of the distribution of the estimators of the bounds, where  $N$  is the number of auctions observed; all of HT’s distribution function bounds are transformations of distribution functions of observables. We start with the estimation of the ME value *density* function  $f^*$ , which is piecewise constant since  $\mathcal{F}^*$  is piecewise linear.

We obtain an estimator of  $f^*$  that is at worst  $\sqrt{N}$ -consistent: if no constraints are binding in the ‘population’ then convergence is arbitrarily fast, otherwise the rate is  $\sqrt{N}$ . However, even under  $\sqrt{N}$ -convergence asymptotic normality does not obtain even if the bound estimates were asymptotically normal since the limit distribution depends on which constraints are binding in the



population.<sup>15</sup> We propose a constraint selection procedure which resembles the moment selection procedure of Andrews and Soares (2010, AS) *inter multa alia*, but differs in three important respects: (a) in AS uniform inference is an important objective but our goal is to derive an expression for the limit distribution of the estimator of  $f^*$ ; (b) AS is intended for the case of set identified parameters whereas  $f^*$  is unique; (c) in AS moment equalities and inequalities are all information available, whereas here we have an optimization problem with a known concave objective function with estimated inequality constraints. Like AS, our constraint selection procedure requires a sample–size–dependent input parameter. We pair the constraint selection procedure with a simulation method to obtain a procedure that is (pointwise) asymptotically similar for some value distributions  $\mathcal{F}^*$  and conservative for others. There is nothing that makes our estimation problem unique to auction environments or indeed to the maximum entropy problem, so our new theoretical results should have wider application. For example, in a game–theoretic discrete choice model where moment inequalities are available to partially identify payoff parameters, our inference methods can be used to construct a confidence interval for the payoff for each player when the only information available is that the payoff parameters satisfy the moment inequalities. For a nonzero length identified set, this problem translates into two separate constrained optimization problems: one for the minimum and one for the maximum.

We further obtain results for the optimal reserve price and the corresponding expected revenue. Since the expected revenue function corresponding to the ME value distribution  $\mathcal{F}^*$  need not be strictly pseudoconcave, the ME reserve price may not be unique. However, we show that for  $\mathcal{F}^*$  the set of reserve price values for which expected revenue is maximized is at most finite: each maximizer can be consistently estimated.

Given that there can be multiple expected revenue maximizers, one still needs to pick one. Note that all elements in the solution set yield the same expected revenue for the ME value distribution in contrast to, say, the HT identified set. Therefore, by the entropy criterion, all elements in the solution set are equivalent. For most purposes, simply choosing a reserve price that maximizes the *estimated* expected revenue will be satisfactory. Doing so has the advantage that one does not have to estimate the entire set of maximizers, which entails the choice of an input parameter: this is our recommended approach. If one insists on estimating the entire set then one can introduce tie breakers. For example, one could compare the worst case loss for each maximizer. A more sophisticated alternative approach is to search for a value density that maximizes entropy among the distributions that are orthogonal to the original ME density  $f^*$ . Finally, one can simply focus on the smallest element of the set of ME optimal reserve price values. This approach can be justified from a welfare perspective.<sup>16</sup>

If one intends to construct a confidence interval for the optimal reserve price or maximum

---

<sup>15</sup>The bounds estimators are not generally asymptotically normal, either, because they are minima of several possibly asymptotically normal quantities.

<sup>16</sup>The seller is indifferent and the buyer gains if the probability of a sale increases with the same expected revenue.

attainable expected revenue for the ME distribution then our recommended approach will not suffice. Hence, for the purpose of inference on the optimal reserve price, we focus our efforts on the simplest approach, i.e. choosing the smallest element in the set of ME optimal reserve price values. We show that our estimator of the smallest ME optimal reserve price has similar statistical properties to our estimator of  $f^*$ : a convergence rate no worse than  $\sqrt{N}$  and a limit distribution that is not normal, but can be simulated. Our estimator of expected revenue, however, is  $\sqrt{N}$ -consistent in all cases. For expected revenue a simulation-based method can be used to conduct inference, also.

Our paper is organized as follows. In section 2 we set up the environment. In section 3, we derive the formulation of the ME solution of  $f^*$ . In section 4 we use examples to compare HT's, AK's, and our approaches. Section 5 contains an empirical illustration. Then in section 6 we develop statistical properties of our estimator of  $f^*$ , the ME optimal reserve price and expected revenue. Section 6 also contains a modest simulation study.

## 2. Preliminaries

As mentioned in section 1, we consider an English auction with symmetric bidders and exogenous entry under the IPV paradigm, in which there is a minimum bid increment  $\Delta > 0$ . Unlike HT, we assume that bid increments are multiples of  $\Delta$ , i.e. one of  $\Delta, 2\Delta, \dots$ , albeit that any other known discrete scheme works, also.<sup>17</sup> In what follows, we shall refer to HT as HT's approach with this additional assumption. Thus, bidders' values  $v_1, \dots, v_n$  are independent and identically distributed (i.i.d.) draws from an unknown continuous distribution function  $\mathcal{F}$  with positive density function  $f$ . We assume that the potential number of bidders  $n$  is known. Throughout the paper we assume that the support of  $v_i$  is the unit interval  $[0, 1]$ . The text is phrased as though we observe the highest bid of each bidder, but observing the winning bid (and the number of potential bidders) is sufficient if one sets the unobserved bids to zero, albeit that the distribution function bounds are then wider. Because this is an English auction, there is no one-to-one correspondence between the observed bid and a bidder's value. Further, since  $\Delta > 0$ , the bid distribution is discrete.

The objective in the empirical auctions literature is typically to uncover  $\mathcal{F}$  from the bid distribution, which is then used to obtain policy-relevant objects such as the expected revenue and optimal reserve price. However, as HT point out, in our setup (point) identification does not obtain. One reason is that  $\Delta > 0$ . If  $\Delta$  were equal to zero then  $\mathcal{F}$  can be point identified under plausible assumptions such as the absence of jump bidding. If  $\Delta$  is nonzero then the bid distributions provide bounds on  $\mathcal{F}$ . Below we briefly review HT's results and issues surrounding them.

It is well-known from the order statistics literature that if  $u_1, \dots, u_n$  are independent with standard uniform distributions then  $u_{i:n}$ , the  $i^{\text{th}}$  (smallest) order statistic, has a beta distribution with parameters  $i$  and  $n - i + 1$ , i.e.  $u_{i:n} \sim \mathcal{B}(i, n - i + 1)$ . Consequently,  $\mathcal{F}(v_{i:n})$  has a  $\mathcal{B}(i, n - i + 1)$

<sup>17</sup>For instance, small increments at low bid levels and large increments at high bid levels.

distribution whose distribution function will be denoted by  $\mathcal{H}_{i:n}$ . Let  $\mathcal{G}_{i:n}$  be the distribution function of the  $i^{\text{th}}$  lowest bid  $b_{i:n}$  and let  $r \in [0, 1]$  be the reserve price.

**Theorem 1** (Haile and Tamer, 2003). Suppose that bidders do not bid more than they are willing to pay and that they do not allow an opponent to win at a price that they are willing to beat. Then, for all  $v \in [r, 1]$ , we have  $\mathcal{F}_L(v) \leq \mathcal{F}(v) \leq \mathcal{F}_U(v)$ , where

$$\mathcal{F}_L(v) = \mathcal{H}_{n-1:n}^{-1}\{\mathcal{G}_{n:n}(v - \Delta)\} \quad \text{and} \quad \mathcal{F}_U(v) = \min_{i=1,2,\dots,n} \mathcal{H}_{i:n}^{-1}\{\mathcal{G}_{i:n}(v)\}. \quad \square$$

There are a few comments to make. First, to facilitate the discussion we assume that the number of participants is constant across auctions: an extension to different numbers of bidders is simple but messy. Second, the bounds in theorem 1 can be estimated. Indeed, given a sample of size  $N$ ,  $\sqrt{N}$ -consistent estimators of  $\mathcal{F}_L$ ,  $\mathcal{F}_U$  can be constructed. Further, the bounds  $\mathcal{F}_L$ ,  $\mathcal{F}_U$  are step functions because the support of  $b_i$  is discrete since  $\Delta > 0$ . Finally, the bounds on  $\mathcal{F}$  do not say much about the density function  $f = \mathcal{F}'$ , which is needed to analyze e.g. the optimal reserve price. In this section we will be mostly concerned with the density issue.

We now turn to the analysis of the optimal reserve price in a counterfactual environment in which  $\Delta = 0$  and the assumptions for the revenue equivalence theorem in Myerson (1981) are satisfied. This is essentially the same exercise as in HT, albeit that we do *not* impose HT's pseudoconcavity assumption. As noted in section 1,<sup>18</sup> jump bidding is effectively ruled out in this counterfactual environment. Let  $\tilde{\pi}(r; \mathcal{F})$  be the expected revenue function in this thought experiment, i.e.

$$\begin{aligned} \tilde{\pi}(r, \mathcal{F}) &= \mathbb{E}\{\max(v_{n-1:n}, r)\mathbb{1}(v_{n:n} > r)\} \\ &= 1 - r\mathcal{F}^n(r) + \int_r^1 \{(n-1)\mathcal{F}^n(v) - n\mathcal{F}^{n-1}(v)\} dv. \quad (1) \end{aligned}$$

So, like HT and AK, we use data on auctions with  $\Delta > 0$  to study the optimal reserve price in regular second price auctions, i.e. second price auctions with  $\Delta = 0$  in which the assumptions for the revenue equivalence theorem are satisfied. This is clearly not ideal but, without further assumptions about bidder behavior, it is the best that can be done. Indeed, with only the minimal behavioral assumptions made in HT, AK, and here, the conditions for Myerson (1981)'s revenue equivalence theorem are not met: for instance, the assumptions are not sufficient for the existence of Nash equilibrium bidding strategies in English auctions.<sup>19</sup> Conversely, if the assumptions necessary for the revenue equivalence theorem were satisfied for the auctions in the data then point identification

<sup>18</sup>See footnote 4.

<sup>19</sup>To establish lemma 4 in their paper, HT make additional assumptions, including the feasible auction mechanism requirement we mentioned in the introduction.

could obtain.<sup>20</sup>

Thus, the optimal reserve price  $r_0$  satisfies

$$\frac{\partial_r \tilde{\pi}(r_0, \mathcal{F})}{n \mathcal{F}^{n-1}(r_0)} = 1 - \mathcal{F}(r_0) - r_0 f(r_0) = 0, \quad (2)$$

provided that  $r_0 \in (0, 1)$ .

Solving the first order condition in (2) requires knowledge about the density function  $f$ . Even then, the solution need not be unique. Therefore, the bounds in theorem 1 generally do not provide much, if any, information about  $r_0$ .

HT's approach in this situation is to note that the right-hand side of (2) is the derivative of  $\mathcal{Q}(r) = r\{1 - \mathcal{F}(r)\}$  at  $r_0$ , and they restrict the function  $\mathcal{Q}$  to be strictly pseudoconcave. This extra assumption on the shape of  $\mathcal{Q}$  ensures that  $r_0$  is uniquely defined and it allows them to derive bounds on  $r_0$  from the bounds on  $\mathcal{F}$ . This solution is defensible because similar assumptions have been made in the auctions literature. But there are limitations: strict pseudoconcavity of  $\mathcal{Q}$  rules out some plausible value distributions. For instance, the distribution described in example 1 has a unimodal density that appears to be perfectly 'regular' yet violates the strict pseudoconcavity assumption.

**Example 1.** Suppose that  $f(v) = 1/(4v^2)$  for  $v \in (1/2, \eta)$  with  $\eta < 1$  and that  $f$  is unrestricted on  $[0, 1/2] \cup [\eta, 1]$  except that  $\mathcal{F}(1/2) = 1/2$ . Then, for any  $v \in (1/2, \eta)$ , we have  $\mathcal{F}(v) = 1 - 1/4v$  so that  $\mathcal{Q}'(v) = 0$ . Therefore,  $\mathcal{Q}$  is not strictly pseudoconcave.  $\square$

The density  $f$  in example 1 satisfies neither Myerson (1981, page 66)'s condition nor HT's. But Myerson shows that his condition is not needed for the characterization of an optimal auction mechanism: it is simply helpful for the computation of one. In section 4 we provide more examples using mixtures of two Beta distributions, where the corresponding function  $\mathcal{Q}$  has two distinct local maxima and hence is not pseudoconcave.

Recall that instead of using HT's assumption that restricts the set of admissible value distributions, we apply the maximum entropy principle. Letting  $\mathcal{F}^*$  be the entropy-maximizing value distribution, our approach is to maximize  $\tilde{\pi}(r, \mathcal{F}^*)$ , whereas AK's proposal is to maximize  $\tilde{\pi}(r, \mathcal{F}_U)$ . To highlight the difference between the two, suppose that the reserve price used in the data is  $\bar{r} > 0$ . Then, all bidders with values less than  $\bar{r}$  (and some with values greater than  $r$ ) will not bid, i.e. they will bid zero.<sup>21</sup> Therefore,  $\mathcal{F}_U(r)$  is flat on  $[0, \bar{r})$ , and it will be positive when the price distribution has positive probability mass at zero. But then (1) shows that no point in  $[0, \bar{r})$  can be the optimal reserve price suggested by AK, unless  $\mathcal{F}_U(r) = 1$  for all  $r$ , since  $\tilde{\pi}(r, \mathcal{F}_U)$  is increasing in  $r \in [0, \bar{r})$ .

<sup>20</sup>The requirements for Myerson's revenue equivalence theorem are not met and point identification does not obtain for several reasons. For starters, one cannot identify a continuous value distribution from a discrete bid distribution. Further, if  $\Delta > 0$  then the winning bidder can end up paying more or less than the value of the second highest bidder: the sequence in which bids are submitted matters. If the winning bidder wins with bid  $\bar{b}$  then that only means that the second highest bidder's value is in the interval  $[\bar{b} - \Delta, \bar{b} + \Delta)$ . (This is not intended to be a complete list.)

<sup>21</sup>Bidders with values greater than zero may not bid because others have already bid past their value.

So AK's optimal reserve price cannot be less than  $\bar{r}$ . This is not the case with the entropy approach. As will become clear in section 3,  $\mathcal{F}^*$  is linear on  $[0, \bar{r}]$ . Therefore, (1) shows that if  $\Omega = \mathcal{F}_U(0)$  is large enough then the maximum entropy optimal reserve price is  $\bar{r} / 2\Omega$ . To put that into perspective suppose that in 9,999 out of 10,000 observed auctions in the data with two bidders and a reserve price equal to 1/2 there are no positive bids. So,  $\hat{\mathcal{G}}_{1:2}(r) \geq 0.9999$  and  $\hat{\mathcal{G}}_{2:2}(r) = 0.9999$  for all  $r \in [0, 1/2)$ . Therefore,  $\hat{\mathcal{F}}_U(r)$  equals either 0.99 or  $\sqrt{0.9999}$  on  $[0, 1/2)$ . Then AK would suggest to keep the reserve price at 1/2 and our maximum entropy solution would suggest dropping the reserve price to approximately 1/4.<sup>22</sup> Keeping the reserve price at 1/2 would be optimal if the value distribution indeed has a large mass point at zero (or something extremely close to it), but in most other cases dropping the reserve price would be better.

### 3. Maximum entropy density

All we know about the value distribution is that it satisfies the bounds in theorem 1. Thus, the maximum entropy density, i.e. the least informative density given the information provided by the identified bounds is given by

$$f^* = \operatorname{argmin}_f \int_0^1 f(s) \log f(s) \, ds \quad \text{subject to} \quad \begin{cases} \int_0^1 f(s) \, ds = 1, \\ \forall v : \mathcal{F}_L(v) \leq \int_0^v f(s) \, ds \leq \mathcal{F}_U(v), \end{cases} \quad (3)$$

where the bounds  $\mathcal{F}_L, \mathcal{F}_U$  were given in theorem 1 and  $\mathcal{F}(v) = \int_0^v f(s) \, ds$  is continuous in  $v$ .

The optimization problem in (3) is simpler than a typical infinite-dimensional optimization problem because the bounds  $\mathcal{F}_L$  and  $\mathcal{F}_U$  are step functions, as noted earlier. Below we reformulate (3) as a finite-dimensional convex optimization problem.

Suppose that for some  $J \geq 2$  the support of the bid distribution consists of the points  $0 = \beta_0 < \beta_1 < \dots < \beta_J < \beta_{J+1} = 1$ . For the sake of presentational simplicity, we assume that the  $\beta_j$ 's are equally spaced, i.e.  $\beta_j = j\Delta$ .

**Lemma 3.1.** The solution  $f^*$  to (3) is constant on each  $I_j = [\beta_{j-1}, \beta_j)$  for  $j = 1, 2, \dots, J+1$ .  $\square$

Therefore, solving (3) is a finite-dimensional problem in terms of its complexity. Specifically, (3) can be solved by finding

$$g_1^*, \dots, g_{J+1}^* := \operatorname{argmin}_{g_1, \dots, g_{J+1} \geq 0} \sum_{j=1}^{J+1} g_j (\log g_j - \log \Delta)$$

<sup>22</sup>Here, we assume that it is known that the support of the value distribution starts at zero. We thank an anonymous referee for pointing this out.

$$\text{subject to } \begin{cases} \sum_{j=1}^{J+1} g_j = 1, \\ \Lambda_{0j} \leq \sum_{k=1}^j g_k \leq \Upsilon_{0j} \quad \text{for } j = 1, 2, \dots, J, \end{cases} \quad (4)$$

where  $g_j = \Delta f(\beta_{j-1})$  and  $0 \leq \Lambda_{0j} := \mathcal{F}_L(\beta_j) \leq \Upsilon_{0j} := \mathcal{F}_U(\beta_{j-1}) \leq 1$ .<sup>23</sup> Since  $g_1^*, \dots, g_{J+1}^*$  sum to one, we define  $g^* = [g_1^*, \dots, g_J^*]^\top \in \mathbb{R}^J$ , omitting  $g_{J+1}^*$ . We show in appendix C that the sign constraints are never binding, i.e.  $g_j^* > 0$  for all  $j$ .

The bounds  $\Lambda_0 = [\Lambda_{01}, \dots, \Lambda_{0J}]^\top$  and  $\Upsilon_0 = [\Upsilon_{01}, \dots, \Upsilon_{0J}]^\top$  are unknown but can be estimated at the parametric rate. Estimation and inference will be discussed later. The vector of bounds will be denoted by  $D_0 = [\Lambda_0^\top, \Upsilon_0^\top]^\top$ . When we wish to emphasize the dependence of  $g^*$  on  $D_0$  we will use the function  $g^\circ$  to write  $g^* = g^\circ(D_0)$ .

The optimization problem in (4) is a finite-dimensional convex programming problem, for which many well-known algorithms are available (e.g. Bertsekas, 2015). In fact, since the objective function in (4) is strictly convex (e.g. by the log sum inequality) and all constraints are linear, the solution to (4) is unique.

The maximum entropy density function  $f^*$  can be obtained from  $g^* = g^\circ(D_0)$ :

$$f^*(v) = f^*\{v, g^\circ(D_0)\} = \sum_{j=1}^{J+1} g_j^\circ(D_0) \mathbb{1}(v \in I_j) / \Delta, \quad (5)$$

where  $\mathbb{1}$  is the indicator function. Consequently, the maximum entropy distribution function  $\mathcal{F}^*$  is given by

$$\mathcal{F}^*(v) = \mathcal{F}^*\{v, g^\circ(D_0)\} = \int_0^v f^*\{s, g^\circ(D_0)\} ds = \sum_{j=1}^{J+1} \mathbb{1}(v \in I_j) a_j^\top(v) g^\circ(D_0), \quad (6)$$

where  $a_j(v) = [1, 1, \dots, 1, v/\Delta - j + 1, 0, 0, \dots, 0]^\top$ .<sup>24</sup> Figure 2 shows an example using simulated bids.

For figure 2, the bounds are estimated with  $S = 100,000$  simulated auctions with  $n \in \{2, 6\}$  potential bidders,  $\Delta = 0.1$ , and two different value distributions. Note that the bounds depend on the bidding strategies used. Here, bids were generated by randomly choosing among the currently losing bidders whose values were at least equal to the current bid level plus  $\Delta$  and assign a bid equal to the current bid level plus  $\Delta$  to the chosen bidder.

<sup>23</sup>The lower bound in (4) comes from the fact that  $\mathcal{F}$  is known to be continuous such that  $\lim_{v \uparrow \beta_j} \mathcal{F}(v) = \mathcal{F}(\beta_j) \geq \mathcal{F}_L(\beta_j)$ . For details, see the proof of lemma 3.1 in appendix C.

<sup>24</sup>For  $v \in I_j = [\beta_{j-1}, \beta_j) = \Delta[j-1, j)$ , we have  $\mathcal{F}^*(v) = a_j^\top(v) g^* = \sum_{k=1}^{j-1} g_k^* + (v/\Delta - j + 1) g_j^*$ .




Figure 2: The HT bounds and the maximum entropy solution: the top figures are based on 100,000 simulated auctions with  $n = 2$ ,  $\Delta = 0.10$  using Beta(0.5, 0.5) (left) and Beta(2, 2) (right) as the true value distributions. The bottom figures use  $n = 6$  with the other settings the same.

When  $n$  is larger, the lower bounds tend to be zero for a wide range of small values of  $v$ , which is because the empirical distribution function of the largest bid tends to be zero for “small” values of  $v$  with finite  $S$ . This becomes an undesirable issue for the upper bound: the upper bound estimated by the usual empirical distribution function will be zero for “small” values of  $v$ . In order to avoid this problem, we used  $\hat{\mathcal{G}}_{i:n}(v) = \{\sum_{s=1}^S \mathbb{1}(b_{i:n,s} \leq v) + 1\} / (S + 1)$ , where  $b_{i:n,s}$  is the  $i^{\text{th}}$  (smallest) bid in auction  $s$ . Please note that  $\hat{\mathcal{G}}_{i:n}$  is asymptotically equivalent to the usual empirical distribution but is always positive with finite  $S$ .

As figure 2 shows, the maximum entropy density generally differs from the true value density. This is not surprising since the true value density is not point-identified. We propose the maximum entropy density as a representative value distribution under partial identification: by the maximum entropy principle it is the least informative choice among the value distributions in the identified set.

Now consider the maximum entropy expected revenue function,  $\pi(r, g^*) = \tilde{\pi}\{r, \mathcal{F}^*(\cdot, g^*)\}$ . The maximizer and the maximum of  $\pi(\cdot, g^*)$  are the maximum entropy optimal reserve price and the corresponding expected revenue, respectively. As we mentioned in section 1, the maximum entropy optimal reserve price is always contained in HT’s interval for the optimal reserve price. However, the maximum entropy distribution itself need not satisfy HT’s strict pseudoconcavity assumption.

## 4. Comparison of methods

We now compare the three approaches to the seller’s problem, namely the bounding approach of Haile and Tamer, the maxmin approach of Aryal and Kim, and our entropy–based approach using some examples. We draw values independently from distributions chosen by us and simulate bids, for which we adopt the algorithm described in HT’s example 2 in their appendix B. We then compute the identified bounds of the value distribution function using  $S = 100,000$  auctions to simulate the objects of estimation, i.e. the true identified bounds. The bounds on the value distribution function are then used to find the (bounds on the) reserve prices proposed by HT, AK, and the present paper.<sup>25</sup> Therefore, the focus of this section is on making a comparison of the different approaches at the population level; this way we highlight the fundamental differences of the different approaches. In the discussion below, for a given value distribution function  $\mathcal{F}$ , we shall refer to the function  $\tilde{\pi}(r, \mathcal{F})$  defined in (1) as the profit function and to  $\mathcal{Q}(r) = r\{1 - \mathcal{F}(r)\}$  as the pseudoprofit function.

In our experiments we use  $n = 2$  since the reserve price has the greatest impact when the number of bidders is small: there is sufficient competition when  $n$  is large. Note that  $n$  here is the number of bidders in the hypothetical second price auction: the number of bidders in the data is immaterial for the purposes of the present exercise. The results described are based on  $\Delta = 0.10$  unless otherwise indicated.

For figure 3 we used Beta(0.5, 0.5) and Beta(2, 2) as the value distributions, resulting in strictly pseudoconcave pseudoprofit functions, as HT requires. Figure 3 has four panels: for each of these two distributions there is one graph for the pseudoprofit function and one for the profit function. In each graph we draw the ‘truth,’ the (pseudo)profit function corresponding to the maximum entropy solution and its maximizer (marked ME), the HT bounds, and the maxmin optimal reserve price, marked AK. The punkish nature of the lower and upper bounds is due to the fact that the bounds on the distribution function are step functions: see figure 2.<sup>26</sup> The maximum entropy solution is a value at which the maximum entropy profit function is maximized which, in the case of strict pseudoconcavity, coincides with the point at which the maximum entropy pseudoprofit function is maximized. The maxmin solution is the point at which the lower bound to the profit function

<sup>25</sup>So, what is marked as AK uses all bids, as we mentioned in section 1.

<sup>26</sup>HT choose to smooth out the bounds, but there is no information on the bounds between nodes of the distribution.






Figure 3: Pseudoprofit (left) and profit (right) for the Beta(0.5, 0.5) (top) and Beta(2, 2) (bottom) value distributions;  $n = 2$ ,  $\Delta = 0.1$ . Green represents the design and black the maximum entropy solution. Upper and Lower show the (pseudo)profits corresponding to the upper and lower bound of the value distribution function, respectively.

is maximal, which is always at one (or more) of the nodes. Finally, to obtain the HT bounds one takes the AK solution and extends it left and right to the point at which the upper bound to the pseudoprofit function attains the maximum of the lower bound.

The HT bounds for the optimal reserve price are well defined and informative in both designs depicted in figure 3, albeit that they are too wide to be of much use. In the Beta(0.5, 0.5) design, maxmin yields a higher profit whereas in the Beta(2, 2) design maximum entropy does better.<sup>27</sup> Also, note that the HT bounds in both designs contain the maximum entropy solution. This phenomenon arises because of the continuity of the maximum entropy distribution function. Since the pseudoprofit function corresponding to the maximum entropy distribution is continuous, its maximum value can

<sup>27</sup>The green profit curve is higher at the value marked AK than at the value marked ME in the top graph and lower in the bottom one.

never be smaller than the maximum of the worst case pseudoprofit function.




Figure 4: The top graphs show a density similar to the one discussed in example 1, along with the corresponding distribution function (green), its bounds (orange and blue), and the maximum entropy solution (black). The bottom figures show the pseudoprofit function (left) and the profit function (right), where the blue and orange lines correspond to the lower and upper bound of the value distribution function presented in the top right panel.

Figures 4 to 6 illustrate what can happen if HT’s strict pseudoconcavity assumption is violated. First, in figure 4 the pseudoprofit function has a flat area; see example 1. In this example, the support of the value distribution is larger than  $[0, 1]$ ; we only draw the unit interval because that is where the action is. We set  $f(v) = -12v^2 + 8v$  on  $[0, 0.5)$  and  $f(v) = 1/(4v^2)$  on  $[0.5, 1)$ , with the remainder of the mass at or beyond 1. In this example, the HT identified set is still convex.

Second, figures 5 and 6 are based on a mixture of two Beta distributions, where the corresponding pseudoprofit functions have two distinct local maxima. Because HT’s assumptions are violated, their results do not apply here. However, applying HT’s machinery to the bounds of the pseudoprofit function produces a nonconvex set for the optimal reserve price. Neither the entropy approach




Figure 5: The top graphs shows the mixture density  $0.9 \cdot \text{Beta}(2, 10) + 0.1 \cdot \text{Beta}(20, 2)$  and the corresponding distribution function (green), along with its bounds (orange and blue), and the maximum entropy solution (black). The bottom figures show the pseudoprofit function (left) and the profit function (right), where the blue and orange lines correspond to the lower and upper bound of the value distribution function presented in the top right panel.

nor the maxmin solution requires the pseudoconcavity assumption to yield a point decision for the reserve price.

Figure 6 shows that the maximum entropy reserve price and the AK solution can be substantially different. In this example, the profit generated by maximum entropy is considerably higher than that generated by maxmin. However, this is not always true, as can be seen in figure 5.

Figure 7 depicts results for a uniform value distribution with  $\Delta = 0.10$  and  $\Delta = 0.20$ . In this case the entropy-maximizing distribution is uniform, also, so the true pseudoprofit and profit functions coincide with their maximum entropy counterparts. There are a few points to note here. First, the HT bounds become less informative as  $\Delta$  increases: for  $\Delta = 0.20$ , one end point of the HT bounds is uninformative in this example. Second, in contrast to maximum entropy, AK always take




Figure 6: The top graphs shows the mixture density  $0.92 \cdot \text{Beta}(2, 10) + 0.08 \cdot \text{Beta}(20, 2)$  and the corresponding distribution function (green), along with its bounds (orange and blue), and the maximum entropy solution (black). The bottom figures show the pseudoprofit function (left) and the profit function (right), where the blue and orange lines correspond to the lower and upper bound of the value distribution function presented in the top right panel.

a value from the mass points of the bid distribution. For instance, if  $\Delta = 0.20$  then the maximum entropy optimal reserve price equals 0.5, which is not in the support of the bid distribution.




Figure 7: The true value distribution is uniform on the unit interval. The top graphs are the pseudoprofit (Left) and the profit function (Right) with  $\Delta = 0.10$ . The bottom graphs use  $\Delta = 0.20$ . The blue and orange lines correspond to the lower and upper bound of the value distribution function.

## 5. Empirical illustration

We now use a modest empirical example to illustrate our method and to make a comparison with the other approaches. We follow HT in using timber auction data and we refer the reader to HT for a more detailed discussion of the features of that data set.

The application is imperfect in that the minimum bid increments in these data are small (in the order of \$0.05 per MBF<sup>28</sup>), with the consequence that the theoretical upper and lower bounds are close and their estimates can cross.

We use HT's method of smoothing the estimates of the lower and upper bounds, denoting

---

<sup>28</sup>Thousand board feet.

the smoothing parameter  $\rho_{HT}$ . We focus here on auctions with two or three bidders to reduce heterogeneity. Like HT, we use auction data from region 6 and zone 2 in the period 1982–1989, where we focus on English auctions that employed the flat pricing method in which the object was sold; this left us with 2,020 auctions to use for this exercise. We incorporate the seller’s value  $v_0$  into the profit function. Since the choices made are not identical to those in HT, our numbers do not coincide with theirs.

We control for (observed) heterogeneity by conditioning on (logarithms of) manufacturing costs, logging costs, selling value, six-month inventory, a measure of species concentration and year dummies; all dollar values are deflated to October 1, 1982 using the producer price index (WPU085). Below we refer to these covariates as  $x$ . Summary statistics for the covariates (except the year dummies) are presented in table 1.

	Mean	Std Dev	Min	Max
mfg costs	209.4	35.1	40.0	294.8
logging costs	133.4	33.2	48.5	358.4
selling value	451.5	64.9	243.4	1303.2
six-month inventory	1653.0	545.0	56.8	2862.4
species concentration	0.7	0.2	0.2	1.0

Table 1: Observed auction heterogeneity

To address the curse of dimensionality we assume that the joint distribution of the valuation and the number of participants given the vector  $x$  of covariates depends on  $x$  only through the single index  $x^\top \beta$ . We estimate  $\beta$  by using a semiparametric regression of the number of participants on covariates using Ichimura’s estimator (e.g. Ichimura, 1993; Hardle, Hall, and Ichimura, 1993). Like HT, we focus on the mean values of the covariates. Thus, to start we estimate bounds for  $\mathcal{F}(v \mid x^\top \beta = \mu_x^\top \beta)$ , for which we use bounds on  $\mathcal{F}(v \mid n = n, x^\top \beta = \mu_x^\top \beta)$  together with  $\mathbb{P}(n = n \mid x^\top \beta = \mu_x^\top \beta)$ : independence between  $v$  and  $n$  (given covariates) is *not* assumed here.

We present our results for  $\rho_{HT} \in \{-3, 0, 3\}$ ; the larger one chooses  $\rho_{HT}$ , the more conservative are the bounds. Figure 8 shows the estimated bounds and the resulting ME solution for the case in which the covariates are evaluated at their means. Since  $\Delta$  is small, there are now many nodes. For  $\rho_{HT} = -3$  the lower and upper bounds almost coincide, and they often cross, which suggests that we almost have point identification. For  $\rho_{HT} = 0, -3$ , the ME solution hugs the lower bound which contrasts with AK’s maxmin approach which is based on the upper bound of the value distribution (which corresponds to the lower envelope of the (pseudo-) profit function).

The recommended optimal reserve prices<sup>29</sup> for various values of  $v_0$  are presented in table 2. As expected, the solutions are far apart for  $\rho_{HT} = 0$  or 3. However, for  $\rho_{HT} = -3$ , the bounds on

<sup>29</sup>In HT’s case bounds on the optimal reserve price.




Figure 8: Estimated bounds of the value distribution

the value distribution are so tight that they lead to some unexpected numbers in table 2. This can be explained by the shape of the pseudo-profit functions corresponding to the bounds of the value distribution, as depicted in figure 9.

$v_0$	$\rho_{HT}$	Entropy	maxmin	$r_L$	$r_U$	$(r_L + r_U)/2$
	-3	\$ 108.82	\$ 129.98	\$ 100.97*	\$ 144.38*	\$ 122.67*
\$20	0	\$ 121.00	\$ 108.82	\$ 72.94	\$ 212.81	\$ 142.88
	3	\$ 121.00	\$ 100.97	\$ 57.34	\$ 237.58	\$ 147.46
	-3	\$ 137.53	\$ 136.16	\$ 129.95*	\$ 205.18*	\$ 167.57*
\$40	0	\$ 132.40	\$ 136.16	\$ 87.00	\$ 212.81	\$ 149.91
	3	\$ 132.40	\$ 137.53	\$ 72.94	\$ 237.58	\$ 155.26
	-3	\$ 204.86	\$ 187.56	\$ 136.16*	\$ 208.09*	\$ 172.12*
\$60	0	\$ 143.89	\$ 136.16	\$ 102.06	\$ 237.58	\$ 169.82
	3	\$ 143.89	\$ 204.86	\$ 88.23	\$ 237.58	\$ 162.90
	-3	\$ 204.86	\$ 187.56	\$ 187.56	\$ 212.81	\$ 200.19
\$80	0	\$ 187.56	\$ 187.56	\$ 125.74	\$ 237.58	\$ 181.66
	3	\$ 152.37	\$ 204.86	\$ 110.82	\$ 237.58	\$ 174.20

\* indicates that the identified set is not convex.

Table 2: Entropy, HT, and AK solutions of the reserve price

For instance, consider the case of  $\rho_{HT} = -3$  and  $v_0 = \$40$ . Here, the upper and lower envelopes of the pseudo-profit functions are not strictly pseudoconcave. When there are two local maximizers (with the local maxima close to each other), our approach picks the smaller one; we did not make this adjustment for the AK solutions reported here. Note that the HT identified set is not convex in this case, which suggests that the value distribution itself may not be strictly pseudoconcave.




Figure 9: Pseudo profit functions for various values of  $v_0$



## 6. Distribution theory

So far we have treated  $D_0$  as known or given. In the remainder we address the fact that  $D_0$  is estimated in practice and develop a distribution theory for the maximum entropy solution, the optimal reserve price and maximum expected revenue. Our problem is similar to that of Hsieh, Shi, and Shum (2018) in that we deal with a constrained optimization problem, but ours is neither linear nor quadratic programming.

We return to the homogeneous auctions case. Our first assumption pertains to the asymptotic behavior of  $\hat{D}$ .

**Assumption A.** For some random vector  $\Phi$ ,  $\hat{\Omega} = \sqrt{N}(\hat{D} - D_0) \xrightarrow{d} \Phi$  as  $N \rightarrow \infty$ .

Since  $D_0$  is a finite-dimensional vector of bounds (see theorem 1) which are probabilities, constructing a  $\sqrt{N}$ -consistent estimator  $\hat{D}$  is a routine exercise. However, without using HT-type smoothing across bidders, the limit distribution  $\Phi$  is not usually normal because of the minimum function that appears in the upper bound; one can however simulate from  $\Phi$ .

Equation (4) has a unique solution  $g^*$  once  $D_0$  is given; recall that we wrote  $g^* = g^\circ(D_0)$  to make this dependence explicit in our discussion below (4). Since  $g^*$  is the unique solution to (4), it follows from the maximum theorem that  $g^\circ$  is continuous. Therefore, consistency of  $\hat{g}^*$  follows from the continuous mapping theorem. Formal proofs of all results are provided in the appendices.

**Theorem 2.** Suppose that assumption A holds. Then  $\hat{g}^* - g^* = o_p(1)$ . □

The maximum entropy solution based on  $\hat{D}$  may not yield a unique optimal reserve price but, as will become apparent in section 6.3, the set of optimal reserve prices is finite. If one only desires to know *an* optimal reserve price then choosing an element from  $\mathcal{R}(\hat{g}^*) = \operatorname{argmax}_r \pi(r, \hat{g}^*)$  will do. If only the optimal profit is desired then  $\mathcal{P}(\hat{g}^*) = \max_r \pi(r, \hat{g}^*)$  is a consistent estimator. For a full discussion of inference-related issues, we refer to section 6.3.

We now develop a distribution theory for ME solutions followed by some Monte Carlo simulations.

**6.1 Asymptotic distribution for  $\hat{g}^*$ :** We start with  $\hat{g}^*$ . The theory developed here is not specific to the entropy problem but can be useful more generally for optimization problems with estimated inequality constraints.

We first explore how perturbations to  $D_0$  affect  $g^*$ . The function  $g^\circ$  is not generally differentiable at  $D_0$ . However, in lemma D.2 we show that it is directionally differentiable in every direction, which can be used to obtain results for the asymptotic distribution of  $\sqrt{N}(\hat{g}^* - g^*)$ . In order to describe the directional derivative, we need to discuss the behavior of the Lagrange multipliers of the optimization problem in (4).

## 6.1 Asymptotic distribution for $\hat{g}^*$

We first eliminate  $g_{J+1}$  and one equality constraint by replacing  $g_{J+1}$  with  $1 - \sum_{j=1}^J g_j$ . The objective function becomes

$$Q(g) = \sum_{j=1}^J g_j (\log g_j - \log \Delta) + \left(1 - \sum_{j=1}^J g_j\right) \left\{ \log \left(1 - \sum_{j=1}^J g_j\right) - \log \Delta \right\} \quad (7)$$

with constraints

$$G_j \leq \Upsilon_{0j} \quad \text{and} \quad G_j \geq \Lambda_{0j} \quad \text{for } j = 1, 2, \dots, J, \quad (8)$$

where  $G_j = \sum_{k=1}^j g_k$ . We define  $G_j^* = \sum_{k=1}^j g_k^*$ . Let  $\lambda_{uj}^* \geq 0$  and  $\lambda_{lj}^* \geq 0$  be the Lagrange multipliers corresponding to the  $\Upsilon_{0j}$  and  $\Lambda_{0j}$  constraints, respectively. Further, let  $\gamma_j^* = \lambda_{uj}^* - \lambda_{lj}^*$ , where we note that  $\lambda_{uj}^* \lambda_{lj}^* = 0$  for all  $j$  because  $\Upsilon_{0j} \geq \Lambda_{0j}$ .<sup>30</sup> The bounds  $D_0 = [\Upsilon_0^\top, \Lambda_0^\top]^\top$  belong to the parameter space  $[0, 1]^{2J}$ . Below, we partition  $[0, 1]^{2J}$  into finitely many areas in such a way that the signs of all the  $\gamma_j^*$ 's are the same for two points in the same area and at least one multiplier has a different sign for two points in different areas; this partition is unique. Thus, each area corresponds to a set of nonzero Lagrange multipliers.

Let  $\gamma^* = [\gamma_1^*, \dots, \gamma_J^*]^\top = \gamma^\circ(D_0)$  for some  $\gamma^\circ$ . Further, let  $K = (K_u, K_\ell)$  be a pair of disjoint sets such that  $K_u \cup K_\ell \subseteq \{1, 2, \dots, J\}$  and define

$$S_K = \left\{ D \in [0, 1]^{2J} : K_+(D) = K_u, K_-(D) = K_\ell, \forall j : \Lambda_j \leq G_j^\circ(D) \leq \Upsilon_j \right\}, \quad (9)$$

where  $K_+(D) = \{j : \gamma_j^\circ(D) > 0\}$ ,  $K_-(D) = \{j : \gamma_j^\circ(D) < 0\}$ , and  $G_j^\circ = \sum_{k=1}^j g_k^\circ$ . So,  $K_u, K_\ell$  represent binding upper and lower bounds, respectively, where we define *binding constraints* to be constraints whose Lagrange multipliers are nonzero; Lagrange multipliers can equal zero even if constraints hold with equality.<sup>31</sup> This is an important distinction as will become apparent in section 6.2.

The  $S_K$  sets are distinct and form a partition of  $[0, 1]^{2J}$  by construction. Consider the following example.

**Example 2.** Suppose that  $J = 2$  and there are only upper bound constraints: i.e.

$$\begin{aligned} & \min_{g_1, g_2, g_3} \{g_1 \log g_1 + g_2 \log g_2 + g_3 \log g_3 - (g_1 + g_2 + g_3) \log \Delta\} \\ & \text{subject to} \begin{cases} g_1 + g_2 + g_3 = 1, \\ g_1 \leq \Upsilon_{10}, \\ g_1 + g_2 \leq \Upsilon_{20}. \end{cases} \end{aligned}$$

<sup>30</sup>It would be more precise to say that there exist solutions for which  $\lambda_{uj}^* \lambda_{lj}^* = 0$  because the Lagrange multipliers are not unique when  $\Upsilon_{0j} = \Lambda_{0j}$ .

<sup>31</sup>For instance, if one minimizes  $x^2$  subject to  $x \leq 0$  then the Lagrange multiplier equals zero but the constraint holds with equality, i.e.  $x = 0$ .

### 6.1 Asymptotic distribution for $\hat{g}^*$

We first eliminate  $g_3$  by using the equality constraint and focus on  $g_1, g_2$ . Then, the Karush–Kuhn–Tucker (KKT) conditions are

$$\begin{cases} \log g_1^* - \log(1 - g_1^* - g_2^*) + \gamma_1^* + \gamma_2^* = 0, & \gamma_1^*(g_1^* - \Upsilon_{10}) = 0, & g_1^* \leq \Upsilon_{10}, \\ \log g_2^* - \log(1 - g_1^* - g_2^*) + \gamma_2^* = 0, & \gamma_2^*(g_1^* + g_2^* - \Upsilon_{20}) = 0, & g_1^* + g_2^* \leq \Upsilon_{20}, \\ \gamma_1^* \geq 0, & \gamma_2^* \geq 0. \end{cases}$$

So, there are four cases:<sup>32</sup>  $(\gamma_1^*, \gamma_2^*) \in \{(0, 0), (+, 0), (0, +), (+, +)\}$ . Each case represents a

$(\gamma_1^*, \gamma_2^*)$	$g_1^*$	$g_2^*$	$D_0 = (\Upsilon_{10}, \Upsilon_{20}) \in$
(0, 0)	1/3	1/3	$S_{(\emptyset, \emptyset)} = \{3\Upsilon_1 \geq 1, 3\Upsilon_2 \geq 2\}$
(+, 0)	$\Upsilon_{10}$	$(1 - \Upsilon_{10})/2$	$S_{(\{1\}, \emptyset)} = \{3\Upsilon_1 < 1, \Upsilon_1 + 1 \leq 2\Upsilon_2\}$
(0, +)	$\Upsilon_{20}/2$	$\Upsilon_{20}/2$	$S_{(\emptyset, \{2\})} = \{3\Upsilon_2 < 2, \Upsilon_2 \leq 2\Upsilon_1\}$
(+, +)	$\Upsilon_{10}$	$\Upsilon_{20} - \Upsilon_{10}$	$S_{(\{1, 2\}, \emptyset)} = \{2\Upsilon_1 < \Upsilon_2, 2\Upsilon_2 < \Upsilon_1 + 1\}$

Table 3: Solutions for the case  $J = 2$  with no lower bounds

unique set of constraints with nonzero multipliers and corresponds to a polygon in  $[0, 1]^2$  as shown in figure 10. In table 3 each such polygon is denoted by  $S_{(K_u, \emptyset)}$  for some  $K_u$ .<sup>33</sup> If two polygons share a boundary then the boundary belongs to the polygon with more multipliers equal to zero.  $\square$




Figure 10: Graphical illustration of the  $S_K$  sets in table 3;  $(0, 0)$ ,  $(+, 0)$ ,  $(+, +)$ , and  $(0, +)$  indicate the corresponding signs of the multipliers  $(\gamma_1^*, \gamma_2^*)$ .

In example 2, the partitioning sets  $S_K$  are polygons. As we show in lemma C.1 in appendix C, the  $S_K$  sets are always polyhedra, which is important for the directional differentiability of  $g^\circ$ . Further,

<sup>32</sup>For general  $J$  and with both upper and lower bounds, there are  $\sum_{r=0}^J 2^r \binom{J}{r}$  different cases.

<sup>33</sup>Since there are no lower bound constraints in this example,  $K_\ell = \emptyset$ .

### 6.1 Asymptotic distribution for $\hat{g}^*$

it is generally true that boundaries belong to the area with more multipliers equal to zero, a property that will prove useful for developing an inference procedure.

Recall that each of the  $S_K$  sets corresponds to a set of *binding constraints*. Therefore, if  $D_0 \in S_K$  then the relevant constraints in (8) can be expressed as

$$R_K^\top g = D_{0K}, \quad (10)$$

where  $R_K \in \mathbb{R}^{J \times \Xi}$  matrix and  $D_{0K} = [\Upsilon_{0K_u}^\top, \Lambda_{0K_\ell}^\top]^\top \in \mathbb{R}^\Xi$  with  $\Xi$  the cardinality of  $K_u \cup K_\ell$ . Here,  $\Upsilon_{0K_u}$  and  $\Lambda_{0K_\ell}$  are the subvectors of  $\Upsilon_0$  and  $\Lambda_0$  determined by the sets  $K_u, K_\ell$  of indices, respectively: for a vector  $D = [\Upsilon^\top, \Lambda^\top]^\top \in \mathbb{R}^{2J}$  and a set of indices  $K = (K_u, K_\ell)$ , the operation of finding the subvector  $D_K = [\Upsilon_{K_u}^\top, \Lambda_{K_\ell}^\top]^\top$  will be denoted by the function  $\phi$  (i.e.  $D_K = \phi(D, K)$ ). Note that  $R_K$  is a matrix of full column rank consisting of zeros and ones, unless  $K_u = K_\ell = \emptyset$ .<sup>34</sup>

**Example 3.** Again consider example 2, where there are four  $S_K$  sets. Since there are no lower bound constraints, we have  $K_\ell = \emptyset$  in all four cases.

(a)  $K = (\emptyset, \emptyset)$ :  $R_K$  is void and no constraints are relevant.

(b)  $K = (\{1\}, \emptyset)$ :  $R_K = [1, 0]$  and  $D_{0K} = \Upsilon_{10}$ .

(c)  $K = (\{2\}, \emptyset)$ :  $R_K = [1, 1]$  and  $D_{0K} = \Upsilon_{20}$ .

(d)  $K = (\{1, 2\}, \emptyset)$ :  $R_K = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$  and  $D_{0K} = \begin{bmatrix} \Upsilon_{10} \\ \Upsilon_{20} \end{bmatrix}$ . □

We now derive an asymptotic expansion of  $\hat{Z} = \sqrt{N}(\hat{g}^* - g^*)$ , which will be the basis for our discussion in section 6.2. Let  $K_0 = (K_{u0}, K_{\ell0})$ , where

$$K_{u0} = \{j : \gamma_j^\circ(D_0) > 0\} \quad \text{and} \quad K_{\ell0} = \{j : \gamma_j^\circ(D_0) < 0\}. \quad (11)$$

So,  $D_0 \in S_{K_0}$  by definition. Define  $\hat{K}$  by replacing  $D_0$  in the definition of  $K_0$  with  $\hat{D}$ , so  $\hat{D} \in S_{\hat{K}}$  by definition as well. Further, let  $\Theta(d, K) = H^{-1} R_K (R_K^\top H^{-1} R_K)^{-1} \phi(d, K)$  when  $K \neq (\emptyset, \emptyset)$  and  $\Theta(d, K) = 0$  when  $K = (\emptyset, \emptyset)$ , where  $H$  is the Hessian of  $Q$  evaluated at  $g^*$ .<sup>35</sup> We then have the following theorem.

**Theorem 3.** Suppose that assumption A holds. Then

$$\hat{Z} := \sqrt{N}(\hat{g}^* - g^*) = \begin{cases} \Theta(\hat{\Phi}, K_0) + o_p(1), & \text{if } D_0 \in \text{int}(S_{K_0}), \\ \Theta(\hat{\Omega}, \hat{K}) + o_p(1), & \text{if } D_0 \in \text{bdr}(S_{K_0}), \end{cases}$$

where  $H$  is the Hessian of  $Q$  at  $g^*$ . □

<sup>34</sup>If  $K_u = K_\ell = \emptyset$  then  $R_K$  is void and all constraints evaporate.

<sup>35</sup>Note that  $H^{-1} = \text{Diag}(g^*) + g^* g^{*\top}$ .

## 6.2 Inference on $g^*$

Please note that theorem 3 does by itself not provide guidance for inference. For instance, we do not know whether or not  $D_0 \in \text{int}(S_{K_0})$ . If  $D_0$  is a boundary point of  $S_{K_0}$ , then  $\hat{K}$  is random and depends on  $\hat{\Omega}$  even in the limit.<sup>36</sup> We propose an inference procedure and establish its validity in section 6.2.

**6.2 Inference on  $g^*$ :** Our inference procedure detects automatically whether  $D_0$  is an interior point or not. If  $D_0$  is an interior point then our procedure is (asymptotically) similar, whereas if  $D_0$  is a boundary point then our procedure is conservative. Our bound selection procedure is similar to what is used in the moment inequality literature (e.g. Andrews and Soares, 2010), but our inference procedure is different. Indeed, unlike Andrews and Soares (2010) we have an optimization problem with inequality constraints with estimated bounds that has a unique solution. Here, the optimization problem stems from our use of maximum entropy. The following example illustrates the basic idea.

**Example 4.** Consider  $\min_{g_1} \{g_1 \log g_1 + (1 - g_1) \log(1 - g_1)\}$  s.t.  $g_1 \leq U_{10}$ . Here, the KKT conditions are given by  $\log g_1^* - \log(1 - g_1^*) + \gamma_1^* = 0$ ,  $\gamma_1^*(g_1^* - U_{10}) = 0$ ,  $\gamma_1^* \geq 0$ . Now, let  $S_0 = \{U : U > 1/2\}$ ,  $S_1 = \{U : U < 1/2\}$ , and  $S_b = \{1/2\}$ . Let  $\hat{U}_1$  be a  $\sqrt{N}$ -consistent estimator of  $U_{10}$ .

- (1) If  $U_{10} \in S_0$ , then  $g_1^* = 1/2$  and  $\gamma_1^* = 0$ ; the constraint holds with (strict) inequality. Further, since  $\hat{U}_1 \in S_0$  with probability approaching one, we have  $\hat{g}_1 = 1/2$  with probability approaching one. Therefore,  $\sqrt{N}(\hat{g}_1^* - g_1^*) \simeq 0$ .
- (2) If  $U_{10} \in S_1$ , then  $g_1^* = U_{10}$  and  $\gamma_1^* = \log\{(1 - U_{10})/U_{10}\} > 0$ ; the constraint holds with equality. Since  $\hat{U}_1 \in S_1$  with probability approaching one, we have  $\hat{g}_1 = \hat{U}_1$ , which means that  $\sqrt{N}(\hat{g}_1^* - g_1^*) \simeq \sqrt{N}(\hat{U}_1 - U_{10})$  in this case.
- (3) If  $U_{10} \in S_b$ , then  $g_1^* = 1/2$  and  $\gamma_1^* = 0$ ; the constraint holds with equality. In this case, the distribution of  $\hat{g}_1$  depends on whether  $\hat{U}_1 \in S_0$  or  $\hat{U}_1 \in S_1$ . That is, if  $\hat{U}_1 \downarrow 1/2$ , then  $\hat{g}_1 = 1/2$  with probability approaching one, while if  $\hat{U}_1 \uparrow 1/2$ , then  $\hat{g}_1 - g_1^* = \hat{U}_1 - U_{10}$  converges at the rate of  $\sqrt{N}$ . Therefore,  $\sqrt{N}(\hat{g}_1 - g_1^*) \simeq \sqrt{N}(\hat{U}_1 - U_{10})\mathbb{1}(\hat{U}_1 > U_{10})$ .

In both cases (2) and (3) the constraint holds with equality but the two cases should be distinguished because the limit distribution of  $\hat{g}_1$  is different: the critical value from case (2) is too small for case (3). Using the value of the multiplier together with the constraint itself can help us to distinguish between the three cases.

In this simple example the limit distribution in case (3) can be obtained explicitly, which is not true in a more general setup. However, we can always take a conservative approach in the boundary case. For instance, in case (3), we can use the fact that  $\max\{0, \sqrt{N}(\hat{U}_1 - U_{10})\} \geq \sqrt{N}(\hat{U}_1 - U_{10})\mathbb{1}(\hat{U}_1 > U_{10}) \simeq \sqrt{N}(\hat{g}_1^* - g_1^*)$ .  $\square$

<sup>36</sup>For instance, suppose that in example 2,  $D_0 = (\Upsilon_{10}, \Upsilon_{20}) = (1/3, 1/3) \in S_{(\emptyset, \emptyset)}$ . Hence,  $\gamma_1^*(D_0) = 0$ . However, for any small  $t > 0$ ,  $\{d : \gamma_1^*(D_0 + td) = 0, \|d\| = 1\}$  and  $\{d : \gamma_1^*(D_0 + td) > 0, \|d\| = 1\}$  are continua.

## 6.2 Inference on $g^*$

Below we elaborate on this idea. Recall from (11) that  $K_0 = (K_{u0}, K_{\ell0})$  are the sets of indices of the constraints that have nonzero multipliers at  $D_0$ . Recall further that we refer to  $K_0$  as the set of *binding* constraints at  $D_0$ , but that there can be constraints that hold with equality that are not in  $K_0$ .

Indeed, define  $K_0^* = (K_{u0}^*, K_{\ell0}^*)$  with

$$K_{u0}^* = \{j : G_j^\circ(D_0) = \Upsilon_{0j}\} \quad \text{and} \quad K_{\ell0}^* = \{j : G_j^\circ(D_0) = \Lambda_{0j}\}. \quad (12)$$

We then have the following lemma.

**Lemma 6.1.**  $K_0 = K_0^*$  if and only if  $D_0 \in \text{int}(S_{K_0})$ .<sup>37</sup> □

Lemma 6.1 is a consequence of the continuity of the solution and the multipliers. The proof is provided in appendix G.

Below, we develop estimators  $\tilde{K}$  and  $\tilde{K}^*$  such that  $\mathbb{P}(\tilde{K} \neq K_0) = o(1)$ ,  $\mathbb{P}(\tilde{K}^* \neq K_0^*) = o(1)$ , and  $\mathbb{P}(\tilde{K} \subseteq \hat{K} \subseteq \tilde{K}^*) = 1$ .<sup>38</sup> In view of theorem 3 and lemma 6.1, we propose simulating the distribution of  $\hat{T} = T(\tilde{K}, \tilde{K}^*)$  for given values of  $\tilde{K}$  and  $\tilde{K}^*$ , where

$$T(K^\circ, K^*) = \max_{K^\circ \subseteq K \subseteq K^*} \Theta(\Phi^*, K), \quad (13)$$

with  $\Phi^*$  an independent copy of  $\Phi$ ; since  $\Theta$  is a vector-valued function, the maximum is element-wise.<sup>39</sup> So in each replication we are using a different draw  $\Phi^*$  but the same estimates  $\tilde{K}, \tilde{K}^*$  based on the original data. Let  $T = T(K_0, K_0^*)$ . Since  $\tilde{K} = K_0$  and  $\tilde{K}^* = K_0^*$  with probability approaching one,  $\hat{T} = T$  with probability approaching one, also, so the distinction between  $\hat{T}$  and  $T$  is moot for our asymptotic analysis. The quantiles of  $T$  provide an upper bound for the corresponding quantiles of  $\hat{Z}$ . Further, if  $D_0$  is an interior point then  $K_0 = K_0^*$  and hence  $T = \Theta(\Phi^*, K_0)$  has the same distribution as  $\Theta(\Phi, K_0) = \hat{Z} + o_p(1)$ . Hence, the quantiles of  $T$  and  $\hat{Z}$  coincide in the limit and inference is asymptotically similar.

Let  $\tilde{K} = (\tilde{K}_u, \tilde{K}_\ell)$  and  $\tilde{K}^* = (\tilde{K}_u^*, \tilde{K}_\ell^*)$ , where

$$\begin{cases} \tilde{K}_u = \{j : \gamma_j^\circ(\hat{D}) \geq \kappa_N\}, & \tilde{K}_\ell = \{j : \gamma_j^\circ(\hat{D}) \leq -\kappa_N\}, \\ \tilde{K}_u^* = \{j : G_j^\circ(\hat{D}) \geq \hat{\Upsilon}_j - \kappa_N\}, & \tilde{K}_\ell^* = \{j : G_j^\circ(\hat{D}) \leq \hat{\Lambda}_j + \kappa_N\}, \end{cases}$$

with  $0 < \kappa_N = o(1)$  and  $1 = o(\kappa_N \sqrt{N})$ . Input parameters like  $\kappa_N$  are much discussed in the moment inequality literature; the BIC choice  $\kappa_N^2 = \log N / N$  is popular.<sup>40</sup>

<sup>37</sup>By the KKT condition, we always have  $K_0 \subseteq K_0^*$ . So,  $K_0 \subsetneq K_0^*$  if and only if  $D_0 \in \text{bdr}(S_{K_0})$ .

<sup>38</sup>For pairs of sets  $K = (K_u, K_\ell)$  and  $K^* = (K_u^*, K_\ell^*)$ , we write  $K \subseteq K^*$  when the inclusion holds elementwise.

<sup>39</sup>One may consider a quadratic form or other scalar-valued transformation of  $\Theta$ . We thank an anonymous referee for making this point.

<sup>40</sup>In practice the standard deviations of  $\hat{\Upsilon}_j$  and  $\hat{\Lambda}_j$  as well as the size of  $\hat{\Upsilon}_j - \hat{\Lambda}_j$  should be taken into account in choosing the input parameter. We thank an anonymous referee for pointing this out.

### 6.3 Inference on expected revenue and optimal reserve price

Let  $\Upsilon^\bullet(D_0) = [\Upsilon_1^\bullet(D_0), \dots, \Upsilon_J^\bullet(D_0)]^\top$  and  $\Lambda^\bullet(D_0) = [\Lambda_1^\bullet(D_0), \dots, \Lambda_J^\bullet(D_0)]^\top$ , where  $\Upsilon_j^\bullet(D_0) = G_j^\circ(D_0) - \Upsilon_{j0}$  and  $\Lambda_j^\bullet(D_0) = G_j^\circ(D_0) - \Lambda_{j0}$ .

**Lemma 6.2.** Suppose that assumption A is satisfied. If  $\gamma^\circ(D_0)$ ,  $\Upsilon^\bullet(D_0)$ , and  $\Lambda^\bullet(D_0)$  do not vary with  $N$ , then (a)  $\mathbb{P}(\tilde{K} \neq K_0) = o(1)$ , (b)  $\mathbb{P}(\tilde{K}^* \neq K_0^*) = o(1)$ . If instead  $\gamma^\circ(D_0)$ ,  $\Upsilon^\bullet(D_0)$ , and  $\Lambda^\bullet(D_0)$  can vary with  $N$  then (a')  $\mathbb{P}(K_0 \notin \tilde{K}) = o(1)$ , (b')  $\mathbb{P}(K_0^* \notin \tilde{K}^*) = o(1)$ . Finally, we always have  $\mathbb{P}(\tilde{K} \subseteq \hat{K} \subseteq \tilde{K}^*) = 1$ .

Lemma 6.2, together with theorem 3, provides the basis for using (13) for inference. Also, lemma 6.2 clarifies in what sense our procedure will be uniformly valid because it generally holds that  $T(K_0, K_0^*) \leq T$  with probability approaching one. Throughout the paper we focus on the pointwise case where  $\gamma^\circ(D_0)$ ,  $\Upsilon^\bullet(D_0)$ , and  $\Lambda^\bullet(D_0)$  are all fixed, but lemma 6.2 shows that our procedure will stay valid (though conservative) in a uniform sense as well.

**Theorem 4.** Suppose that assumption A is satisfied. Then  $\mathbb{P}(\hat{T} \neq T) = o(1)$ . Further, for any  $x \in \mathbb{R}^J$ ,  $\mathbb{P}(\hat{Z} \leq x) \geq \mathbb{P}(T \leq x) + o(1)$ , where the inequality holds with equality whenever  $D_0 \in \text{int}(S_{K_0})$ .  $\square$

**6.3 Inference on expected revenue and optimal reserve price:** We now consider estimation of and inference for the maximum attainable revenue and optimal reserve price corresponding to the maximum entropy solution for the value distribution. We build on our discussion in section 6.2.

Consider

$$\mathcal{P}(g^*) = \max_{r \in [0,1]} \pi(r, g^*) \quad \text{and} \quad \mathcal{R}(g^*) = \operatorname{argmax}_{r \in [0,1]} \pi(r, g^*), \quad (14)$$

where  $\pi(r, g) = \tilde{\pi}\{r, \mathcal{F}^*(\cdot, g)\}$ :  $\tilde{\pi}(r, \mathcal{F})$  and  $\mathcal{F}^*(v, g)$  were introduced in sections 2 and 3, respectively. So,  $\pi(r, g^*)$  is the maximum entropy expected profit function.  $\mathcal{P}(g^*)$  can be estimated by  $\mathcal{P}(\hat{g}^*)$  but  $\mathcal{R}(\hat{g}^*)$  need not be a consistent estimator of  $\mathcal{R}(g^*)$ , albeit that  $\mathcal{R}(\hat{g}^*)$  is contained in  $\mathcal{R}(g^*)$  with probability approaching one. So if the sole objective is to select an optimal reserve price, then selecting an element from  $\mathcal{R}(\hat{g}^*)$  suffices.

Below we construct an estimator of  $\mathcal{R}(g^*)$  and determine its properties. Once we establish the limiting distribution of the estimator  $\hat{r}_1$  of the smallest element in  $\mathcal{R}(g^*)$ ,  $\hat{\mathcal{P}}^* = \pi(\hat{r}_1, \hat{g}^*)$  is an estimator of  $\mathcal{P}(g^*)$  which is more convenient for inference purposes than  $\mathcal{P}(\hat{g}^*)$ .

Although  $\mathcal{R}(g^*)$  need not be a singleton, it is finite. The reason is that, for any  $g > 0$ , the function  $\theta(\cdot, g)$  with

$$\theta(r, g) = \theta^+(r, g) = \frac{\partial_r^+ \tilde{\pi}\{r, \mathcal{F}^*(r, g)\}}{n \{\mathcal{F}^*(r, g)\}^{n-1}}, \quad (15)$$

where  $\partial_r^+$  denotes a right derivative, is piecewise linear in  $r$  with negative slope coefficients.<sup>41</sup>

<sup>41</sup>Note that (2) and (15) are identical except that we now do not implicitly assume the existence of the partial derivative and only focus on distribution functions of the form  $\mathcal{F}^*(\cdot, g)$ .

### 6.3 Inference on expected revenue and optimal reserve price

Indeed, for all  $g$ ,

$$\theta(r, g) = 1 - M_{j-1}(g) + (j-1)g_j - 2g_j r / \Delta \quad \text{if } r \in I_j, \quad (16)$$

where  $M_j(g) = G_j = \sum_{k=1}^j g_k$  and  $M_0(g) = 0$ . Note that  $\theta$  is possibly discontinuous at the  $\beta_j$ 's, i.e. on the support of the bid distribution, because  $\theta(\beta_j, g)$  need not equal  $\theta^-(\beta_j, g)$  for any or all  $j$ , which is defined as  $\theta$  in (16) but with  $\partial_r^+$  replaced with the left derivative  $\partial_r^-$ .

**Lemma 6.3.** For some integer  $1 \leq m < \infty$  and some  $0 < r_1^* < r_2^* < \dots < r_m^* < 1$ ,  $\mathcal{R}(g^*) = \{r_1^*, r_2^*, \dots, r_m^*\}$ . Further, each  $\bar{I}_j = [\beta_{j-1}, \beta_j]$  contains at most one element of  $\mathcal{R}(g^*)$ .  $\square$

We show that both the number and the identity of the  $r_j^*$ -values can be estimated consistently: the rate at which  $m$  is estimated is arbitrarily fast. The proof is simple. Let  $\hat{\mathcal{R}}_N = \{r \in [0, 1] : \pi(r, \hat{g}^*) \geq \mathcal{P}(\hat{g}^*) - \kappa_n\}$ . Since  $\pi(\cdot, \hat{g}^*)$  is continuous,  $\hat{\mathcal{R}}_N$  is a compact subset of  $[0, 1]$  by construction. Consistency is obtained in a similar manner as in Chernozhukov, Hong, and Tamer (2007), albeit that here the identified set is known to be a collection of isolated points.

**Theorem 5.**  $d_H\{\hat{\mathcal{R}}_N, \mathcal{R}(g^*)\} = o_p(1)$ , where  $d_H$  denotes the Hausdorff distance.  $\square$

Note that  $\hat{\mathcal{R}}_N$  is set-valued. We can use it to create point estimates of each of the  $r_j^*$ -values as follows. Let  $\tilde{\kappa}_N = \sqrt[3]{\kappa_N}$ ,  $\hat{\mathcal{R}}_{N,1} = \{r \in \hat{\mathcal{R}}_N : r - \min \hat{\mathcal{R}}_N \leq \tilde{\kappa}_N\}$ ,  $\hat{\mathcal{R}}_{N,2} = \{r \in \hat{\mathcal{R}}_N \setminus \hat{\mathcal{R}}_{N,1} : r - \min(\hat{\mathcal{R}}_N \setminus \hat{\mathcal{R}}_{N,1}) \leq \tilde{\kappa}_N\}$ , etcetera: so,  $\hat{\mathcal{R}}_N = \cup_{k=1}^{\hat{m}} \hat{\mathcal{R}}_{N,k}$  where  $\hat{m}$  is the smallest value of  $\tilde{m}$  for which  $\hat{\mathcal{R}}_{N,\tilde{m}+1}$  is empty. We define our estimator of  $r_k^*$  as

$$\hat{r}_k = \min_{r \in \hat{\mathcal{R}}_{N,k}} \operatorname{argmax} \pi(r, \hat{g}^*).$$

Our definition of  $\hat{r}_k$  allows for the possibility that the maximizer of  $\pi(r, \hat{g}^*)$  in  $\hat{\mathcal{R}}_{N,k}$  is not unique. Consistency of  $\hat{r}_k$  for  $r_k^*$  is straightforward to establish.

**Theorem 6.** We have  $\mathbb{P}(\hat{m} \neq m) = o(1)$ , and for all  $k = 1, \dots, m$ ,  $\hat{r}_k - r_k^* = O_p(\sqrt{\kappa_N})$ .<sup>42</sup>  $\square$

As we will show below, the convergence rate of  $\hat{r}_k$  is in fact better than the rate obtained in theorem 6.

We now turn to the construction of confidence intervals and focus on  $r_1^*$ : the arguments are analogous for  $r_2^*, \dots, r_m^*$  if  $m > 1$ . Note that  $r_1^*$  is the unique maximizer of  $\pi(r, g^*)$  on some compact interval  $A_1$ : such an interval exists by lemma 6.3. Define  $r_1(g) = \min \tilde{R}_1(g)$ , where

$$\tilde{R}_1(g) = \operatorname{argmax}_{r \in A_1} \pi(r, g). \quad (17)$$

<sup>42</sup>If  $\hat{m} < m$  then  $\hat{r}_{\hat{m}+1}, \dots, \hat{r}_m$  can be defined arbitrarily.



### 6.3 Inference on expected revenue and optimal reserve price

So the difference between  $r_1(\hat{g}^*)$  and  $\hat{r}_1$  is that the maximization is conducted over a(n asymptotically) larger set in the former case:  $A_1$  versus  $\hat{\mathcal{R}}_{N,1}$ . However, the difference is minute from a theoretical perspective.

**Lemma 6.4.**  $\mathbb{P}\{r_1(\hat{g}^*) \neq \hat{r}_1\} = o(1)$ . □

Typically,  $\tilde{\mathcal{R}}_1(g)$  is a singleton for all  $g$  in a small enough neighborhood of  $g^*$ , but there is one exceptional case in which  $\tilde{\mathcal{R}}_1(g)$  has two elements for some  $g$  near  $g^*$ .<sup>43</sup>

In view of lemma 6.4 we consider the asymptotic distribution of  $\sqrt{N}\{r_1(\hat{g}^*) - r_1(g^*)\}$ . If the function  $r_1$  is (Hadamard-) differentiable at  $g^*$  then the delta method will apply. Unfortunately, as we commented earlier,  $r_1$  may fail to be Hadamard differentiable. However, even in that case, small perturbations of  $g^*$  have only limited effects on the function  $r_1$ , which is sufficient for our purpose. Below we discuss the Hadamard derivatives of  $r_1$ , which requires us to analyze the first order condition of (17).

Recall that  $\theta(\cdot, g^*)$  is piecewise linear but not necessarily continuous. So we distinguish between the case in which  $r_1(g^*) = r_1^*$  is a continuity point of  $\theta$  and the case in which it is not. Let  $j^*$  be such that  $r_1^* \in I_{j^*}$ .




Figure 11: Graphical illustration of the first order condition for  $r_1^*$

Since  $\theta$  is linear and downward sloping on  $I_{j^*}$ , there are two possibilities, which are illustrated in figure 11. First, if  $r_1^* > \beta_{j^*-1}$  then  $\theta\{r_1(g^*), g^*\} = 0$ , in which case the derivative of  $r_1^*$  at  $g^*$  can be obtained via the implicit function theorem. However, if  $r_1^* = \beta_{j^*-1}$ , then  $r_1^*$  may not be differentiable at  $g^*$  and there are then four separate (sub)cases to consider as shown in table 4.

In appendix H.2 we show that  $r_1$  is Hadamard-differentiable in cases I–IV. Case V is the exceptional case mentioned earlier, but even in case V valid inference for  $r_1^*$  is possible.

<sup>43</sup>The exception arises when  $\tilde{r}_1(g^*) = \beta_{j^*-1}$  for some  $j^*$  and  $\theta^+(\beta_{j^*-1}, g^*) = \theta^-(\beta_{j^*-1}, g^*) = 0$ . Then Hadamard differentiability fails and there can be two maximizers. Absent this technicality,  $r_1(g)$  is the unique maximizer of  $\pi(r, g)$  on  $A_1$  in a neighborhood of  $g^*$ .

### 6.3 Inference on expected revenue and optimal reserve price

case	condition
I	$\beta_{j^*-1} < r_1^* < \beta_{j^*}$ ;
II	$r_1^* = \beta_{j^*-1}$ with $\theta(r_1^*, g^*) < 0 < \theta^-(r_1^*, g^*)$ ;
III	$r_1^* = \beta_{j^*-1}$ with $\theta(r_1^*, g^*) = 0 < \theta^-(r_1^*, g^*)$ ;
IV	$r_1^* = \beta_{j^*-1}$ with $\theta(r_1^*, g^*) < 0 = \theta^-(r_1^*, g^*)$ ;
V	$r_1^* = \beta_{j^*-1}$ with $\theta(r_1^*, g^*) = 0 = \theta^-(r_1^*, g^*)$ .

Table 4: Separate cases for the sensitivity analysis

Let  $\delta, \delta^- \in \mathbb{R}^J$  with

$$\delta_k = -\frac{\Delta}{2g_{j^*}^*} \times \begin{cases} 1, & k < j^*, \\ (1 - G_{j^*-1}^*)/g_{j^*}^*, & k = j^*, \\ 0, & k > j^*. \end{cases} \quad \delta_k^- = -\frac{\Delta}{2g_{j^*-1}^*} \times \begin{cases} 1, & k < j^* - 1, \\ (1 - G_{j^*-2}^*)/g_{j^*-1}^*, & k = j^* - 1, \\ 0, & k > j^* - 1. \end{cases}$$

Further, for  $v \in \mathbb{R}^J$ , define

$$\psi_{g^*}(v) = \begin{cases} v^\top \delta & \text{in case I,} \\ 0 & \text{in case II,} \\ \max(0, v^\top \delta) & \text{in case III,} \\ \min(0, v^\top \delta^-) & \text{in case IV,} \\ \max(|v^\top \delta|, |v^\top \delta^-|) & \text{in case V,} \end{cases} \quad (18)$$

which is the Hadamard derivative of  $r_1$  at  $g^*$  in the direction  $v$  in cases I–IV and an upper bound of the effect of perturbing  $g^*$  on  $r_1$  in case V.

The first case in (18) is the most common: it arises with an interior solution ( $r_1^* > \beta_{j^*-1}$ ), in which case  $\theta(r_1^*, g^*) = 0 = \theta^-(r_1^*, g^*)$ . However, this case is dramatically different from  $\theta(r_1^*, g^*) = 0 = \theta^-(r_1^*, g^*)$  with  $r_1^* = \beta_{j^*-1}$ , i.e. case V. Case II is pictured in the right panel of figure 11: small changes in  $g^*$  do not affect  $r_1^*$ . The remaining two cases in (18) arise if one of the two line segments in the right panel of figure 11 has an endpoint at  $(\beta_{j^*-1}, 0)$ .

We now characterize the asymptotic distribution of  $\hat{t} = \sqrt{N}(\hat{r}_1 - r_1^*)$ . Let  $\Theta$  be the dominant right hand side term in theorem 3, i.e. depending on the location of  $D_0$  either  $\Theta(\Phi, K_0)$  or  $\Theta(\hat{\Omega}, \hat{K})$ .

**Theorem 7.** If assumption A is satisfied then in cases I–IV, we have  $\hat{t} = \psi_{g^*}(\Theta) + o_p(1)$ , and in case V,  $|\hat{t}| \leq \psi_{g^*}(\Theta) + o_p(1)$ .  $\square$

Now consider revenue. Recall that  $\mathcal{P}(g^*) = \pi(r_1^*, g^*)$  is estimated by  $\hat{\mathcal{P}}^* = \pi(\hat{r}_1, \hat{g}^*)$ .

### 6.3 Inference on expected revenue and optimal reserve price

**Theorem 8.** If assumption A is satisfied then  $\sqrt{N}\{\hat{\mathcal{P}}^* - \mathcal{P}(g^*)\} = \partial_g \pi(r_1^*, g^*)^\top \Theta + o_p(1)$ .  $\square$

$\kappa = c\kappa_0$	Beta(0.5,0.5), n=2								
$c = 0.5$	0.0000	0.0000	0.9842	0.9842	0.9842	0.9844	0.9878	0.9884	0.0004
$c = 1.0$	0.0000	0.0000	0.9938	0.9938	0.9938	0.9942	0.9964	0.9878	0.0004
$c = 1.5$	0.0000	0.0000	0.9984	0.9984	0.9984	0.9988	0.9998	0.9856	0.0002
	Beta(0.5,0.5), n=3								
$c = 0.5$	0.0000	0.0000	0.9870	0.9870	0.9870	0.9872	0.9896	0.9742	0.0000
$c = 1.0$	0.0000	0.0000	0.9900	0.9900	0.9900	0.9904	0.9942	0.9668	0.0000
$c = 1.5$	0.0000	0.0000	0.9932	0.9932	0.9932	0.9936	0.9970	0.9566	0.0000
	Beta(1.0,1.0), n=2								
$c = 0.5$	0.0000	0.9962	0.9962	0.9962	0.9962	0.9962	0.9962	0.9962	0.9962
$c = 1.0$	0.0000	0.9962	0.9962	0.9962	0.9962	0.9962	0.9962	0.9962	0.9962
$c = 1.5$	0.0000	0.9962	0.9962	0.9962	0.9962	0.9962	0.9962	0.9962	0.9962
	Beta(1.0,1.0), n=3								
$c = 0.5$	0.0000	0.9854	0.9854	0.9854	0.9854	0.9854	0.9854	0.9854	0.9854
$c = 1.0$	0.0000	0.9850	0.9850	0.9850	0.9850	0.9850	0.9850	0.9850	0.9850
$c = 1.5$	0.0000	0.9856	0.9856	0.9856	0.9856	0.9856	0.9856	0.9856	0.9856
	Beta(2.0,2.0), n=2								
$c = 0.5$	0.0002	0.0292	0.1430	0.9826	0.9706	0.9602	0.9710	0.9744	0.6232
$c = 1.0$	0.0002	0.0210	0.1596	0.9868	0.9750	0.9648	0.9754	0.9800	0.7226
$c = 1.5$	0.0002	0.0160	0.1846	0.9900	0.9780	0.9678	0.9784	0.9844	0.7922
	Beta(2.0,2.0), n=3								
$c = 0.5$	0.0000	0.1172	0.0816	0.9826	0.9828	0.9826	0.9828	0.9868	0.6102
$c = 1.0$	0.0000	0.1090	0.0806	0.9848	0.9850	0.9848	0.9850	0.9914	0.6396
$c = 1.5$	0.0000	0.1030	0.0824	0.9874	0.9876	0.9874	0.9876	0.9940	0.6612

Table 5: Coverage rates at each node with nominal rate 0.95,  $N = 1000$ ; 5000 replications.

Recall that the function  $\psi_{g^*}$  depends on the case. For instance, in case II,  $\hat{r}_1$  is superconsistent. In fact, convergence in this case can be shown to be arbitrarily fast. Therefore, inference on  $r_1$  requires that we know which of the five cases we are in. This is not hard to do because theorem 7 shows that the rate of  $\hat{r}_1$  cannot be worse than  $\sqrt{N}$  and hence we can find out which case is relevant with probability approaching one. Therefore, we can rely on theorem 4 to conduct inference on  $r_1^*$ .

For example, suppose that  $\hat{r}_1 \in (\beta_{\hat{j}^*-1} + \kappa_N, \beta_{\hat{j}^*} - \kappa_N)$  for some  $\hat{j}^*$ . We can then conduct inference for  $r_1^*$  by using the distribution of  $\delta^\top \Theta$  for which theorem 4 can be applied. Inference will be conservative in case V but case V is extreme.

Theorem 8 does not distinguish case V from the other four cases. In fact,  $\hat{g}^*$  is the only relevant object for the limit distribution of  $\pi(\hat{r}_1, \hat{g}^*)$ . This phenomenon is an implication of the envelope

## 6.4 Numerical experiments

theorem. Indeed, in cases I and V, both the right and left derivatives of  $\pi(\cdot, g^*)$  at  $r_1^*$  are zero. In case II, neither of the directional derivatives is zero but  $\sqrt{N}(\hat{r}_1 - r_1^*)$  is asymptotically negligible. Similar arguments apply to cases III and IV, also. Therefore, inference on maximum revenue is straightforward: we can use theorem 4 without knowing which case is relevant.

**6.4 Numerical experiments:** Like in section 4 we simulated bids from English auctions using Beta(0.5, 0.5), Beta(1, 1), and Beta(2, 2) as the distributions of valuations. The true values of the bounds and ME solutions were computed by simulation.

Like in section 4 we estimate the bounds  $D_0$  using empirical distribution functions, where we do not smooth out the minimum function for the upper bound. Since the distribution of  $\Phi$  is generally not normal, we use subsampling to obtain random draws from it.

Tables 5 and 6 show the coverage rates of our 95% confidence intervals for  $g^*$  at each node for  $N = 1000$  with  $n = 2, 3$  and  $\Delta = 0.1$ ; we also tried  $N = 500$  and  $N = 700$  but the results are largely the same. We used  $\kappa = c\kappa_0$  with  $c = 0.5, 1, 1.5$  and  $\kappa_0 = \text{const}\sqrt{\log N/N}$  for every node, where const was chosen by multiplying the maximum gap between the upper and lower bound estimates and the largest standard deviation of (the subsample version of)  $\Phi$ .<sup>44</sup> In these experiments the performance of the procedure is decent, though conservative, except around the two end points.

However, in experiments not reported here, we found that the performance can be poor when the number of bidders is large. This can be explained partly by the fact that the joint distribution of the bounds is poorly estimated when the number of bidders is large. Since the ME estimates can be computed efficiently, it can be beneficial to subsample them given that our theory already shows that the solutions have stable distributional behavior in the limit.

$\kappa = 1 \cdot \kappa_0$	$r_1(g^*)$	$\mathcal{P}(g^*)$
Beta(0.5, 0.5), n=2	0.9974	0.9930
Beta(0.5, 0.5), n=3	0.9956	0.9928
Beta(1, 1), n=2	0.9990	0.9994
Beta(1, 1), n=3	1.0	0.9966
Beta(2, 2), n=2	0.9834	0.9388
Beta(2, 2), n=3	0.9928	0.9830

Table 6: Coverage rates with nominal rate 0.95: N=1000 and 5000 replications.

<sup>44</sup>Checking all subsets between  $\tilde{K}$  and  $\tilde{K}^*$  is NP hard in general, but it is typically feasible if  $\kappa_N$  is relatively small.

# Appendices

## A. Optimal auctions

Myerson (1981)'s strict monotonicity assumption on the so-called virtual valuation function implies the strict pseudoconcavity of the function  $\mathcal{Q}$ . Therefore, Myerson (1981)'s results show that absent strict pseudoconcavity of  $\mathcal{Q}$  a second price auction is not necessarily optimal.

With the maximum entropy value distribution in hand, one can construct a Myerson optimal auction, which can yield higher expected revenue for the seller than a second price auction with a maximum entropy optimal reserve price. Doing so would be impossible if one merely had bounds on the value distribution function.

The optimal auction mechanism works as follows. Let

$$T^*(q) = -F^{-1}(q)(1 - q), \quad q \in [0, 1], \quad (19)$$

and let  $T$  be its *convex hull*, i.e. the maximum convex function below or equal to  $T^*$ . Define

$$C(v) = T'\{F(v)\}.^{45} \quad (20)$$

$C$  is continuous and nondecreasing, which plays a critical role in describing the allocation rule of Myerson's optimal auction. If  $T^*$  were convex already, then  $C$  would be the same as  $C^*$  with  $C^*(v) = v - \{1 - F(v)\} / f(v)$ , which would be increasing.

To see how Myerson's allocation rule can be different from that of a second price auction, suppose that  $n = 2$  and  $v_1 > v_2$ . We continue to assume that the seller values the object at zero. Let  $r_m^*$  denote the smallest value of  $v$  for which  $C(v) = 0$ . Further, define  $\bar{b} = \max[\max\{v : C(v) = C(v_2)\}, r_m^*]$ , and let  $\underline{b}$  be the greater of  $r_m^*$  and  $\min\{v : C(v) = C(v_2)\}$ . If  $v_1 > \bar{b}$  then player 1 wins the auction and pays  $(\bar{b} + \underline{b}) / 2$ . If  $v_1 \leq \bar{b}$  then each player wins the auction with probability  $1 / 2$  and the winner pays  $\underline{b}$ . If  $C$  were strictly convex then  $\underline{b} = \bar{b} = \max(v_2, r_m^*)$  and we would be back in the standard second price auction case.

To illustrate, consider figure 12 which is based on a value distribution that is a mixture of Betas.<sup>46</sup> Here, as is apparent from the dotted lines,  $r_m^* = 0.2$ . If  $v_1 < r_m^*$  then the object is not sold and if  $v_2 < r_m^* < v_1$  then player 1 wins the auction and pays  $r_m^*$ . So suppose that  $v_2 \geq r_m^*$ . If  $v_2$  belongs to an area in which  $C$  is increasing (i.e. not flat) then player 1 wins and pays  $v_2$ . Finally, suppose that  $v_2$  is in an area in which  $C$  is flat, say the large flat segment that extends from about 0.38 to about 0.70. If  $v_1$  is also between 0.38 and 0.70 then each player wins with probability 0.50 and the

<sup>45</sup>We are ignoring the easily addressed nuisance that  $T$  may not be differentiable at a countable set of points.

<sup>46</sup>0.95 times a Beta(2,10) plus 0.05 times a Beta(20,2). The corresponding  $\mathcal{Q}$  function has two distinct local maxima. See section 4.




Figure 12: The functions  $C^*$  (blue dashed) and  $C$  (green solid) for a mixture of Betas.

winner pays 0.38. Otherwise, player 1 wins with certainty and pays 0.54.

As noted, the Myerson mechanism yields an expected profit for the seller that is equal to or exceeds that of a second price auction with an optimally chosen reserve price. However, the difference in expected profit is generally small. For instance, in the example of figure 12 the difference is about 0.001 despite the substantial convexity correction evidenced by the difference between dashed and solid lines in figure 12. This is not entirely surprising in view of Hartline (2016, corollary 5.3), which provides bounds on the gain from an optimal auction compared to a second price one and is a corollary to the Bulow–Klemperer theorem (Bulow and Klemperer, 1996), albeit that the bounds are fairly wide if the number of bidders is small. Further, as noted, it can be cumbersome to implement Myerson’s auction in practice.

We therefore focus on the choice of an optimal reserve price in a second price auction in the remainder of this paper.

## B. Entropy, maxmin, and decision theory

We now provide a decision–theoretic view of the ME solution, which is the focus of this paper. Specifically, we consider choosing the reserve price to maximize minimum expected revenue where the minimum is taken over all distributions which satisfy the bound constraints and are moreover such that their entropy is bounded below by a prespecified value  $\mathcal{E}^*$ . Like in AK, the seller is ambiguity–averse but she now also wants to rule out value distributions that are too crazy. Thus,  $\mathcal{E}^*$  represents a tradeoff between ambiguity–aversion and plausibility. In this framework the AK and ME solutions can be understood as special cases: the AK solution corresponds to the case in which  $\mathcal{E}^* = -\infty$ , whereas the ME solution obtains if  $\mathcal{E}^*$  is the maximum achievable entropy.

The worst case revenue for given  $\mathcal{E}^*$  is

$$\min_{\ell \geq 0} \tilde{\pi}(r, \mathcal{F}) \quad \text{subject to} \quad \begin{cases} -\int_0^1 \ell(s) \log \ell(s) \, ds \geq \mathcal{E}^*, \\ \int_0^1 \ell(s) \, ds = 1, \\ \forall v : \mathcal{F}_L(v) \leq \int_0^v \ell(s) \, ds \leq \mathcal{F}_U(v), \end{cases} \quad (21)$$

where  $\tilde{\pi}$  is given in (1),  $\mathcal{E}^*$  is the minimum desired entropy level, and  $\mathcal{F}_U, \mathcal{F}_L$  are still the identified bounds of the distribution function of valuation. Below we show how to solve (21). We focus on the simplest case, i.e. the case with two bidders: if the number of (independent) bidders is large then choosing the optimal reserve price becomes less important.

The problem in (21) involves infinite-dimensional objects and its solution is not generally closed-form. However, it can always be reformulated as a finite-dimensional parametric problem. To see this, note that  $\mathcal{F}_U, \mathcal{F}_L$  are step functions, so the distribution bound constraints in (21) only contain finitely many restrictions. We can thus write the problem as

$$\min_{\ell \geq 0} \left\{ 2r \bar{\mathcal{F}}(r) - r \bar{\mathcal{F}}^2(r) + \int_r^1 \bar{\mathcal{F}}^2(v) \, dv \right\} \\ \text{subject to} \quad \begin{cases} -\int_0^1 \ell(s) \log \ell(s) \, ds \geq \mathcal{E}^*, \\ \bar{\mathcal{F}}(1) = 0, \\ 1 - \Upsilon_{0j} \leq \bar{\mathcal{F}}(\beta_j) \leq 1 - \Lambda_{0j}, \quad j = 1, 2, \dots, J, \end{cases} \quad (22)$$

where  $\bar{\mathcal{F}}(v) = 1 - \mathcal{F}(v)$ ,  $\Lambda_{0j} = \mathcal{F}_L(\beta_j)$ , and  $\Upsilon_{0j} = \mathcal{F}_U(\beta_{j-1})$  as explained right below (4). We then use the first order condition from variational calculus to obtain the following result. Let  $k_r = \min\{j : \beta_j \geq r\}$ ,  $\bar{Z} = \bar{\mathcal{F}}(r)$ , and  $M_j = \mathcal{F}(\beta_j)$ , where we focus on  $r > 0$ .

**Theorem 9.** The solution  $\mathcal{F}$  to (22) is such that (i)  $\bar{\mathcal{F}}$  is continuous; (ii) for  $v \leq r$ ,  $\bar{\mathcal{F}}$  is piecewise linear; and (iii) for  $v > r$ ,

$$\bar{\mathcal{F}}(v; \alpha, \nu, \omega, \tilde{\beta}) = \begin{cases} -\alpha \omega_j \tan\{(v_j - v)\omega_j\} & \text{for } \max(\beta_{j-1}, r) \leq v < \min(\tilde{\beta}_j, \beta_j) \\ 1 - \Upsilon_{0j} & \text{for } \min(\tilde{\beta}_j, \beta_j) \leq v < \beta_j, \end{cases} \quad (23)$$

for  $\alpha < 0$  and  $k_r \leq j \leq J + 1$ . Further, if neither inequality condition holds with equality at  $\beta_j$  then  $\omega_j = \omega_{j+1}$  and  $\nu_j = \nu_{j+1}$ . Finally, the parameter vectors  $\nu, \omega$ , and  $\tilde{\beta}$  are known functions of  $\bar{Z}, \alpha$ , and the identities of the distribution bound inequalities that hold with equality.  $\square$

The proof of theorem 9 is provided in appendix E. The shape of  $\bar{\mathcal{F}}$  for  $v \leq r$  does not affect expected profit, so for  $v \leq r$  the solution coincides with the ME solution. The parametric function

$\mathcal{F}(\cdot; \alpha, \nu, \omega, \tilde{\beta})$  turns out to be the unique solution to a differential equation implied by the first order condition of (22). Here,  $\alpha$  is the Lagrange multiplier for the entropy constraint, and  $\omega, \nu$  are parameters that show up in the solution to the differential equation. The  $\tilde{\beta}$ -parameters appear when the sign restrictions on  $f$  produce corner solutions. The intuition for this is as follows. As  $\mathcal{E}^* \rightarrow -\infty$ , the solution gets closer to AK's pure maxmin solution, i.e. the upper bound of the value distribution, which is discrete. Unless  $\mathcal{E}^*$  is large negative,  $\tilde{\beta}_j = \beta_j$ .

Theorem 9 says that  $\alpha < 0$  in the solution: the entropy constraint is binding for any finite  $\mathcal{E}^*$ . However, as we will show later in this section, the solution to (21) is continuous in  $\mathcal{E}^*, \Lambda_{j0}, \Upsilon_{0j}$ . As  $\mathcal{E}^* \rightarrow -\infty, \alpha \rightarrow 0$ , eventually resulting in  $\tilde{\beta}_j < \beta_j$ .

It is convenient that the problem can be characterized as an optimization problem in two unknowns if one fixes  $r$  and the identities of the bound constraints that hold with equality. This is explained in the proof in appendix E.

By using theorem 9 we compute the solution to (22) in multiple steps. Fixing the set of constraints that hold with equality and the values of  $\tilde{\mathcal{F}}(r)$  and the Lagrange multiplier for the entropy constraint, we solve the parametric problem, which we then minimize over all candidate values of  $\tilde{\mathcal{F}}(r)$  and the entropy constraint Lagrange multiplier. For each set of constraints we thus have a solution. Finally, we minimize over all finitely many binding constraint sets.

We have implemented our method in three designs. The results of our efforts are depicted in figures 13 to 16: Figures 13, 15 and 16 depict the distribution function for which expected profit is minimized for the entropy-constrained maxmin solution for  $r$ . Figure 14 depicts the entropy-constrained minimum profit functions as a function of the reserve price.

In all three designs the nodes are at 0, 0.2, . . . , 0.8, 1. The main difference between the designs is in the bounds: in figures 13, 15 and 16 the lower bounds are yellow and the upper bounds are blue. In all cases, the optimal  $r$  was chosen from the grid 0.01, 0.03, . . . , 0.99.

For each design, four different values of  $\mathcal{E}^*$  were used: -10.0, -1.0, -0.1, -0.01 for design 1 and -10.0, -1.0, -0.1, -0.05 for designs 2 and 3. The reason for the discrepancy is that for the bounds used in designs 2 and 3 there exist no distributions that yield an entropy value in excess of  $-0.01$ .

In design 1, the uniform distribution is the maximum entropy solution resulting in an optimal reserve price equal to 0.5. As can be seen in figures 13 and 14, for  $\mathcal{E}^* = -0.01$  our methodology produces a profit function close to the profit function for a uniform value distribution and a distribution function at the optimal reserve price close to a uniform. As  $\mathcal{E}^*$  decreases, however, the distribution function to the right of  $r$  places more weight on values slightly above node values. This is natural, since putting a lot of weight there corresponds to having a pessimistic view of the world. Since the profit is unaffected by the shape of the distribution left of  $r$ , the distribution left of  $r$  is for all values of  $\mathcal{E}^*$  fairly regular.

Now, minimum expected profit as a function of  $r$  for design 1 has an intuitive shape, also. For large negative values of  $\mathcal{E}^*$ , we get the familiar sawtooth pattern with drops at each of the nodes.






Figure 13: The graphs depict the distribution functions of the least favorable distribution for the optimal choice of  $r$  in design 1, where the least favorable distribution is determined by taking the minimum over all distributions whose entropy exceeds a threshold  $\mathcal{E}^*$ : each panel corresponds to a different choice of  $\mathcal{E}^*$ . In all cases the nodes are at  $0, 0.2, \dots, 1.0$ . The graphs were created using a grid of  $r$ -values, namely  $0.01, 0.03, \dots, 0.99$ . The maximal value of  $r$  is indicated by a vertical dotted line.

The drop arises because if the seller sets the reserve price above the node value then she loses revenue from all bidders with values between the node value and  $r$ : for large negative values of  $\mathcal{E}^*$  the distribution of values is concentrated near the node values.

In design 2, the uniform distribution does not satisfy the bound constraints and hence the solution for values of  $\mathcal{E}^*$  close to zero is not a uniform and  $\mathcal{E}^*$  cannot be pushed to 0. Like with design 1, the profit function becomes more sawtoothed as  $\mathcal{E}^*$  decreases because the least favorable distribution




Figure 14: The graphs depict minimum expected profit as a function of  $r$  for three different choices of upper and lower bounds. The minimum at each value of  $r$  is taken over all distributions whose entropy exceeds a threshold  $\mathcal{E}^*$ : each curve in a given panel corresponds to a different choice of  $\mathcal{E}^*$  with greater values of  $\mathcal{E}^*$  resulting in higher expected profit. In all cases the nodes are at  $0, 0.2, \dots, 1.0$ . The graphs were created using a grid of  $r$ -values, namely  $0.01, 0.03, \dots, 0.99$ .

function at each  $r$  gets closer to the unconstrained least favorable distribution. In other words, design 2 is similar in behavior to design 1.

We deliberately chose a fairly extreme case for design 3 with an upper bound that is constant over a large range of values. That said, the solution would be much the same if the upper bound increased only slightly going from one node value to the next. In contrast to designs 1 and 2, the optimal reserve price changes dramatically depending on the choice of  $\mathcal{E}^*$  and so does expected profit. For instance, if the seller sets the optimal reserve price according to  $\mathcal{E}^* = -10$  then her profit




Figure 15: The graphs depict the distribution functions of the least favorable distribution for the optimal choice of  $r$  in design 2, where the least favorable distribution is determined by taking the minimum over all distributions whose entropy exceeds a threshold  $\mathcal{E}^*$ : each panel corresponds to a different choice of  $\mathcal{E}^*$ . In all cases the nodes are at  $0, 0.2, \dots, 1.0$ . The graphs were created using a grid of  $r$ -values, namely  $0.01, 0.03, \dots, 0.99$ . The maximal value of  $r$  is indicated by a vertical dotted line.

is substantially less if the actual value distribution is closer to the maximum entropy distribution. The converse is also true, but the loss is less.

As promised, we conclude by showing that the solution to (22) has some continuity properties. Let  $\tilde{\mathcal{E}}^*(D_0)$  be the maximum entropy level that is attainable for given  $D_0 = [\Lambda_0^\top, \Upsilon_0^\top]^\top$ . Define  $\bar{\mathcal{E}}^* = \inf_{D_0 \in \mathcal{D}} \tilde{\mathcal{E}}^*(D_0)$ , where  $\mathcal{D}$  is an arbitrary neighborhood of  $D_0$ . We denote the solution to (22) by  $\pi^\bullet(r, \mathcal{E}^*, D_0)$ , which is a function on  $[0, 1] \times [\underline{\mathcal{E}}^*, \bar{\mathcal{E}}^*] \times \mathcal{D}$  for  $\underline{\mathcal{E}}^* > -\infty$ .




Figure 16: The graphs depict the distribution functions of the least favorable distribution for the optimal choice of  $r$  in design 3, where the least favorable distribution is determined by taking the minimum over all distributions whose entropy exceeds a threshold  $\mathcal{E}^*$ : each panel corresponds to a different choice of  $\mathcal{E}^*$ . In all cases the nodes are at  $0, 0.2, \dots, 1.0$ . The graphs were created using a grid of  $r$ -values, namely  $0.01, 0.03, \dots, 0.99$ . The maximal value of  $r$  is indicated by a vertical dotted line.

**Theorem 10.** The function  $\pi^\bullet$  is continuous on  $[0, 1] \times [\mathcal{E}^*, \mathcal{E}^*] \times \mathcal{D}$ . Further,  $\tilde{r}(\mathcal{E}^*, D_0) = \operatorname{argmax}_r \pi^\bullet(r, \mathcal{E}^*, D_0)$  is upper-hemicontinuous on  $[\mathcal{E}^*, \bar{\mathcal{E}}^*] \times \mathcal{D}$ .  $\square$

The proof of theorem 10 can be found in appendix F.

In the remainder of the paper we assume that the seller shares our preference for choosing  $\mathcal{E}^*$  maximally, i.e. for maximum entropy.

## C. Lemmas for the solution

**Proof of lemma 3.1:** Fix any  $j^\circ \in \{1, 2, \dots, J+1\}$  and let  $\mathcal{A} = \{\ell : \forall s \notin I_{j^\circ} : \ell(s) = \ell^*(s)\}$ . Then  $f^*$  is not only a solution to (3), but since  $\mathcal{F}_L, \mathcal{F}_U$  are flat on  $I_{j^\circ}$  also to

$$\min_{f \in \mathcal{A}} \int_{\beta_{j^\circ-1}}^{\beta_{j^\circ}} \ell(s) \log \ell(s) \, ds \quad \text{s.t.} \quad \begin{cases} \int_{\beta_{j^\circ-1}}^{\beta_{j^\circ}} \ell(s) \, ds = c_{j^\circ}, \\ \forall v \in I_{j^\circ} : \mathcal{F}_L(\beta_{j^\circ-1}) \leq \underline{\mathbf{S}} + \int_{\beta_{j^\circ-1}}^v \ell(s) \, ds \leq \mathcal{F}_U(\beta_{j^\circ-}), \\ \mathcal{F}_L(\beta_{j^\circ}) \leq \underline{\mathbf{S}} + \int_{\beta_{j^\circ-1}}^{\beta_{j^\circ}} \ell(s) \, ds \leq \mathcal{F}_U(\beta_{j^\circ-}) \end{cases}$$

where  $c_j = \int_{\beta_{j-1}}^{\beta_j} \ell^*(s) \, ds$ ,  $\underline{\mathbf{S}} = \sum_{j=1}^{J^*-1} c_j$ , and  $\mathcal{F}_U(\beta_{j^*-}) = \lim_{v \uparrow \beta_{j^*}} \mathcal{F}_U(v)$ : the last inequality constraint is a condition coming from the fact that  $\mathcal{F}(\cdot)$  is a continuous function.

Because  $\ell^*$  satisfies the inequality constraints, we must have  $\mathcal{F}_L(\beta_{j^\circ-1}) \leq \underline{\mathbf{S}} \leq \underline{\mathbf{S}} + c_{j^\circ} \leq \mathcal{F}_U(\beta_{j^\circ-})$  and  $\mathcal{F}_L(\beta_{j^\circ}) \leq \underline{\mathbf{S}} + c_{j^\circ}$ , and therefore the inequality constraints are redundant. Hence  $f^*$  is constant on  $I_{j^\circ}$ .  $\square$

The Karush–Kuhn–Tucker (KKT) conditions for minimizing (7) subject to (8) are for  $j = 1, \dots, J$  given by

$$\begin{cases} \log g_j^* - \log(1 - G_j^*) + \sum_{k=j}^J (\lambda_{uk}^* - \lambda_{lk}^*) = \lambda_{sj}^*, & (24a) \end{cases}$$

$$\begin{cases} \lambda_{uj}^* (G_j^* - \Upsilon_{0j}) = 0, \quad G_j^* \leq \Upsilon_{0j}, \quad \lambda_{uj}^* \geq 0, & (24b) \end{cases}$$

$$\begin{cases} \lambda_{lj}^* (G_j^* - \Lambda_{0j}) = 0, \quad G_j^* \geq \Lambda_{0j}, \quad \lambda_{lj}^* \geq 0, & (24c) \end{cases}$$

$$\begin{cases} \lambda_{sj}^* g_j^* = 0, \quad g_j^* \geq 0, \quad \lambda_{sj}^* \geq 0. & (24d) \end{cases}$$

Recall that  $\gamma_j^* = \lambda_{uj}^* - \lambda_{lj}^*$ , where  $\lambda_{uj}^* \lambda_{lj}^* = 0$ . Here, if  $g_j^* = 0$ , then the conditions in (24a) and (24d) cannot be satisfied simultaneously. Thus  $g_j^* > 0$  and  $\lambda_{sj}^* = 0$  for all  $j$ .

**Lemma C.1.** The solution  $g^*$  depends on which constraints are binding but is otherwise an affine function of  $\Upsilon_0, \Lambda_0$ . Therefore, each of the  $S_K$  sets defined in (9) is a polyhedron.

*Proof.* Note that  $\lambda_{uj}^* \lambda_{lj}^* = 0$  for all  $j$  because  $\Upsilon_{0j} \geq \Lambda_{0j}$ .<sup>47</sup> So, the conditions in (24a) to (24c) contain  $2J$  unknowns:  $g_1^*, \dots, g_J^*$  and  $\gamma_1^*, \dots, \gamma_J^*$ . We need to solve

$$\begin{cases} \log g_j^* - \log(1 - G_j^*) + \sum_{k=j}^J \gamma_k^* = 0, & (25a) \end{cases}$$

$$\begin{cases} \gamma_j^* (G_j^* - B_{j0}) = 0, & (25b) \end{cases}$$

<sup>47</sup>It would be more precise to say that there exist solutions for which  $\lambda_{uj}^* \lambda_{lj}^* = 0$  because the Lagrange multipliers are not unique when  $\Upsilon_{0j} = \Lambda_{0j}$ .

for  $j = 1, \dots, J$ , where  $B_{j0}$  is either  $\Upsilon_{0j}$  or  $\Lambda_{0j}$ . The conditions in (25a) imply that

$$\gamma_j^* = \log g_{j+1}^* - \log g_j^*, \quad j = 1, 2, \dots, J, \quad (26)$$

where  $g_{J+1}^* = 1 - G_J^*$ . Suppose that there are  $r$  multipliers  $\gamma_j^*$  that equal zero and  $J - r$  that are nonzero. For  $\gamma_j^* \neq 0$ , by (25b) we have  $G_j^* = B_{j0}$ . For  $\gamma_j^* = 0$ , (26) implies that  $g_{j+1}^* = g_j^*$  with  $1 - G_j^* = g_j^*$  as a special case. Therefore,  $g^* = [g_1^*, \dots, g_J^*]^\top$  is the solution to a linear equation system whose right hand side is linear in the  $B_{j0}$ 's.  $\square$

**Lemma C.2.** The solutions  $g^*, \gamma^*$  are continuous functions of  $\Upsilon_0, \Lambda_0$ .

*Proof.* It suffices to show the continuity of  $g^*$ : the continuity of  $\gamma^*$  then follows from (26). The solution  $g^* = [g_1^*, \dots, g_J^*]^\top$  minimizes  $Q$  (defined in (7)) subject to  $g \in \Xi(\Upsilon_0, \Lambda_0)$ , where  $\Xi$  is the correspondence  $\Xi(\Upsilon_0, \Lambda_0) = \{g : \Lambda_{0j} \leq \sum_{k=1}^j g_k \leq \Upsilon_{0j} \text{ for } j = 1, 2, \dots, J\}$ . Since  $Q$  is a continuous function and  $\Xi$  is a continuous correspondence, it follows from the maximum theorem that  $g^* = g^\circ(\Upsilon_0, \Lambda_0)$  is upper hemicontinuous as a correspondence. Further, by the convexity of the problem,  $g^\circ(\Upsilon_0, \Lambda_0)$  is a single element correspondence, i.e. a function, and therefore upper hemicontinuity is equivalent to continuity.  $\square$

**Proof of theorem 2:** Follows from lemma C.2 and the continuous mapping theorem.  $\square$

## D. Sensitivity of the solution

In this section we consider the effect on the solution  $g^*$  of perturbations of  $\Upsilon_0, \Lambda_0$  in a given direction  $d$ . We will use these results for statistical inference.

Recall that  $\mathcal{S} = \{S_K\}$  is a (finite) partition of  $[0, 1]^{2J}$ , where  $S_K$  is defined in (9). As we discussed in section 6, the solution  $g^\circ(D)$  at  $D = [\Upsilon^\top, \Lambda^\top]^\top \in S_K$  can be expressed as

$$g^\circ(D) = \underset{g}{\operatorname{argmin}} Q(g) \quad \text{subject to} \quad R_K^\top g = D_K, \quad (27)$$

where  $R_K$  is a matrix of ones and zeroes with full column rank and  $D_K = [\Upsilon_{K_u}, \Lambda_{K_\ell}]$  is a subvector of  $D$  that is determined by  $K = (K_u, K_\ell)$ ; if none of the constraints in  $S_K$  are binding (i.e.  $K_u = K_\ell = \emptyset$ ) then the restrictions in (27) evaporate.

Now, suppose that we perturb a given  $D_0 = (\Upsilon_0^\top, \Lambda_0^\top)^\top \in S_{K_0}$  in the direction  $d$ , where  $S_{K_0}$  is implicitly defined. So we consider  $D_0 + td$ , where  $d$  is given and  $t > 0$  is small. The most important insight is that for all sufficiently small  $t > 0$ ,  $D_0 + td$  lies within the set  $S_{K_d}$  which only depends on  $d$ . The following lemma formalizes this idea.

**Lemma D.1.** There exist an  $S_{K_d} \in \mathcal{S}$  and an  $\epsilon > 0$  such that  $D_0 + td \in S_{K_d}$  for all  $0 < t < \epsilon$ .

*Proof.* If  $D_0$  is in the interior of  $S_{K_0}$ , then the assertion is true with  $S_{K_d} = S_{K_0}$ . Suppose that  $D_0$  is on the boundary of  $S_{K_0}$ . By lemma C.1 all  $S_K$  sets in  $\mathcal{S}$  are polyhedra, and therefore there are only four possibilities: for a sufficiently small  $\epsilon > 0$ , the (open) line segment  $\{D_0 + td : 0 < t < \epsilon\}$  (i) is a subset of the boundary of  $S_{K_0}$ ; (ii) is a subset of the boundary of some  $S_K \neq S_{K_0}$ ; (iii) belongs to the interior of  $S_{K_0}$ ; (iv) belongs to the interior of some  $S_K \neq S_{K_0}$ .  $\square$

By lemma D.1, for all sufficiently small  $t > 0$ , the solution at  $D_0 + td$  is given by

$$g^\circ(D_0 + td) = \underset{g}{\operatorname{argmin}} Q(g) \quad \text{subject to} \quad R_{K_d}^\top g = D_{0K_d} + td_{K_d}, \quad (28)$$

where  $D_{0K_d}, d_{K_d}$  are the subvectors of  $D_0, d$  corresponding to the indices in  $K_d = (K_{du}, K_{d\ell})$ , as described in the paragraph after (10). The formulation in (28) is convenient for obtaining directional derivatives of  $g^*$ .

Now, note that

$$\lim_{t \downarrow 0} g^\circ(D_0 + td) = \underset{g}{\operatorname{argmin}} Q(g) \quad \text{subject to} \quad R_{K_d}^\top g = D_{0K_d}. \quad (29)$$

Since  $g^\circ$  is continuous by lemma C.2, (29) is equivalent to

$$g^\circ(D_0) = \underset{g}{\operatorname{argmin}} Q(g) \quad \text{subject to} \quad R_{K_0}^\top g = D_{0K_0}, \quad (30)$$

where  $D_{0K_0}$  is a subvector of  $D_0$  defined by  $K_0 = (K_{0u}, K_{0\ell})$  as in (10).<sup>48</sup>

So the directional derivative  $\nabla g^\circ(D_0, d)$  of  $g^*$  at  $D_0$  in the direction  $d$  can be obtained by differentiating (28) with respect to  $t$ .<sup>49</sup> Recall  $H$  be the Hessian of the objective function  $Q$  at  $g^*$ .

**Lemma D.2.** (i) If  $K_d = (\emptyset, \emptyset)$  then  $\nabla g^\circ(D_0, d) = 0$ . (ii) Otherwise,

$$\nabla g^\circ(D_0, d) = H^{-1} R_{K_d} (R_{K_d}^\top H^{-1} R_{K_d})^{-1} d_{K_d}.$$

*Proof.* Since part (i) is trivial, we prove part (ii). Note that  $H$  is positive definite: the typical element  $h_{ij}$  is given by  $h_{ij} = \{\mathbb{1}(i = j) / g_i^*\} + (1 / g_{J+1}^*)$ . The first order conditions of (28) are  $\partial_g Q\{g^\circ(D_0 + td)\} = R_{K_d} \mu(t)$  and  $R_{K_d}^\top g^\circ(D_0 + td) = D_{K_d} + td_{K_d}$ , where  $\mu(t)$  is the vector of the Lagrange multipliers. By differentiating with respect to  $t$  at 0, we obtain  $H \nabla g^\circ(D_0, d) = R_{K_d} \partial_t \mu(0)$  and  $R_{K_d}^\top \nabla g^\circ(D_0, d) = d_{K_d}$ , which implies that  $d_{K_d} = R_{K_d}^\top H^{-1} R_{K_d} \partial_t \mu(0)$ . Therefore, the assertion follows.  $\square$

<sup>48</sup>Although  $S_{K_0}$  and  $S_{K_d}$  need not be the same when  $D_0$  is a boundary point, equivalence of (29) and (30) is intuitive. Indeed, suppose that  $D_0$  is a boundary point of  $S_{K_0}$  and  $D_0 + td \in S_{K_d}$  for all sufficiently small  $t > 0$ , where  $K_0 \neq K_d$ . Now,  $K_0 \subset K_d$  because  $\gamma_j^\circ(\cdot)$  is continuous in view of lemma C.2. Thus, any constraints that are binding at  $D_0 + td$  become *redundant* in the limit.

<sup>49</sup>So, for  $\bar{g}^* \circ (t) = g^\circ(D_0 + td)$ ,  $\nabla g^\circ(D_0, d) = \partial_t \bar{g}^*(0)$ .

**Lemma D.3.**  $\gamma^*$  is directionally differentiable at  $D_0$ . If  $K_d = (\emptyset, \emptyset)$  then  $\nabla \gamma_j^\circ(D_0, d) = 0$ . Otherwise,  $|\nabla \gamma_j^\circ(D_0, d)| \leq \|(R_{K_d}^\top R_{K_d})^{-1} H \nabla g^\circ(D_0, d)\|$ .

*Proof.* By continuity of  $\gamma_j^*$ , there are three relevant cases: (i)  $\gamma_j^\circ(D_0) = 0$  and  $\gamma_j^\circ(D_0 + td) = 0$ ; (ii)  $\gamma_j^\circ(D_0) = 0$  and  $\gamma_j^\circ(D_0 + td) \neq 0$ ; (iii)  $\gamma_j^\circ(D_0) \neq 0$  and  $\gamma_j^\circ(D_0 + td) \neq 0$ . In the first case,  $\nabla \gamma_j^\circ(D_0, d) = 0$ . In the other two cases,  $\nabla \gamma_j^\circ(D_0, d)$  is an element of  $\partial_t \mu(0)$  in the proof of lemma D.2.  $\square$

**Proof of theorem 3:** We first show that  $\hat{Z} = \nabla g^\circ(D_0, \hat{d} / \|\hat{d}\|) \sqrt{N} \|\hat{d}\| + o_p(1)$ , where  $\hat{d} = \hat{D} - D_0$ . Let for any distance  $\rho > 0$  and direction  $a$  with  $\|a\| = 1$ ,

$$\xi_N(a, \rho) = \sqrt{N} \{g^\circ(D_0 + \rho a / \sqrt{N}) - g^\circ(D_0)\} - \nabla g^\circ(D_0, a) \rho.$$

From lemma D.2, we know that  $\xi_N(a, \rho) = o(1)$  for any  $a, \rho$ . Now, for any measure  $\zeta$  and any  $\epsilon > 0$ ,  $\iint \mathbb{1}\{\|\xi_N(a, \rho)\| > \epsilon\} d\zeta(a, \rho) = o(1)$ , by the dominated convergence theorem. Let  $\hat{\rho} = \sqrt{N} \|\hat{d}\|$  and  $\hat{a} = \sqrt{N} \hat{d} / \hat{\rho}$ . Take  $\zeta$  to be the distribution of  $(\hat{a}, \hat{\rho})$ . Therefore, the boundary case is proved since  $\hat{D} \in S_{\hat{K}}$  is always true by definition. For the interior case, it follows from the fact that  $D_0 \in \text{int}(S_{K_0})$  implies  $\hat{D} \in S_{K_0}$  with probability approaching one.  $\square$

## E. Hybrid problem — solution

**Proof of theorem 9:** The first order conditions to (22) are given by

$$2r \mathcal{F}(r) \mathbb{1}(v \leq r) + 2 \int_{\max(v, r)}^1 \bar{\mathcal{F}}(s) ds + \alpha \{\log \ell(v) + 1\} - \sum_{j=1}^J (\tilde{\gamma}_j - \check{\gamma}_j) \mathbb{1}(v \leq \beta_j) + \tilde{\lambda} = 0, \quad (31)$$

and

$$\begin{cases} - \int_0^1 \ell(v) \log \ell(v) dv = \mathcal{E}^*, \\ \tilde{\gamma}_j \left( \int_0^{\beta_j} \ell(s) ds - \Upsilon_{0j} \right) = 0, & j = 1, \dots, J, \\ \check{\gamma}_j \left( \int_0^{\beta_j} \ell(s) ds - \Lambda_{0j} \right) = 0, \end{cases} \quad (32)$$

where  $\tilde{\gamma}_j, \check{\gamma}_j \geq 0$  and  $\alpha < 0$ :  $\alpha$  cannot be zero since then  $\mathcal{F}_U$  minimizes the profit which implies  $\mathcal{E}^* = -\infty$ . Therefore, for  $k = 1, 2, \dots, J$  and  $\beta_{k-1} < v \leq \beta_k$ , we have

$$2r \mathcal{F}(r) \mathbb{1}(v \leq r) + 2 \int_{\max(v, r)}^1 \bar{\mathcal{F}}(s) ds + \alpha \{\log \ell(v) + 1\} - \sum_{j=k}^J (\tilde{\gamma}_j - \check{\gamma}_j) + \tilde{\lambda} = 0. \quad (33)$$



Now, for  $v \leq r$ , the solution  $f$  to (33) is piecewise constant. For  $v > r$ , differentiating (33) yields

$$2\tilde{\mathcal{F}}(v) - \alpha \frac{f'(v)}{f(v)} = 0, \quad (34)$$

whenever  $f(v) > 0$ . Therefore, there is no loss of generality in merging  $(\beta_{k-1}, \beta_k]$  and  $(\beta_k, \beta_{k+1}]$  when the distribution bound constraints are not holding with equality at  $\beta_k$ . Below we focus on  $(\beta_{k-1}, \beta_k]$ , where we have  $\tilde{\gamma}_{k-1} > 0$  and  $\tilde{\gamma}_k > 0$ , i.e. the (upper) bounds at the two end points are binding; binding lower bounds are dealt with analogously.

If  $f(v) = 0$  at any  $v \in (\beta_{k-1}, \beta_k]$  then there must be some  $\tilde{\beta}_k \in (\beta_{k-1}, \beta_k]$  such that  $f(v) = 0$  for all  $\tilde{\beta}_k < v \leq \beta_k$ : putting  $f(v) = 0$  anywhere other than on the right side of  $(\beta_{k-1}, \beta_k]$  would not minimize profit. So (34) must be satisfied for all  $\beta_{k-1} < v \leq \tilde{\beta}_k$ . The proposed parametric form of  $f$  is a solution to the second order differential equation in (34).

To see that the final statement of the theorem is true consider the following. By the definition of  $k_r$  we start with  $3(J + 2 - k_r)$  unknowns: one set  $\{\omega_j, \nu_j, \tilde{\beta}_j\}$  for each interval. We have  $\tilde{\mathcal{F}}(r) = \bar{Z}$  and  $\tilde{\mathcal{F}}(1) = 0$  as two binding equalities. Suppose that there are no binding constraints at  $\beta_{k_r}, \beta_{k_r+1}, \dots, \beta_J$ . Then  $\tilde{\beta}_j = \beta_j$ ,  $\omega_j = \omega_{j+1}$ , and  $\nu_j = \nu_{j+1}$  for all  $j \geq k_r$ . That gives us  $2 + (J + 2 - k_r) + 2(J + 1 - k_r) = 3(J + 2 - k_r)$  equalities, as desired.<sup>50</sup> If a constraint is binding at some  $\beta_j$  then we have two intervals with binding endpoint constraints and the above argument can be repeated for each interval separately. For large negative values of  $\mathcal{E}^*$  it can happen that there are no values of  $\omega_j, \nu_j$  to make both bounds at the endpoints of the interval binding in which case  $\tilde{\beta}_j$  is set to the largest value for which both bounds can be made binding and  $\omega_j, \nu_j$  are then the corresponding solutions.  $\square$

## F. Hybrid problem — continuity

We now turn to the proof of theorem 10. Let  $\ell = 1, \dots, \bar{L}$  denote each of the cases in which the identity of the inequality constraints that hold with equality is given. Define  $\pi_\ell^\circ(\bar{Z}, \alpha, \mathcal{E}^*, D_0; r)$  to be the minimum revenue for a given  $r$  as a function of  $\bar{Z} = \tilde{\mathcal{F}}(r)$ ,  $\alpha$ ,  $\mathcal{E}^*$ , and  $D_0$ . So,

$$\pi^\bullet(\mathcal{E}^*, D_0; r) = \min_{\ell=1,2,\dots,\bar{L}} \left\{ \min_{(\bar{Z}, \alpha) \in \mathcal{C}_\ell(\mathcal{E}^*, D_0)} \pi_\ell^\circ(\bar{Z}, \alpha, \mathcal{E}^*, D_0; r) \right\}, \quad (35)$$

where  $\mathcal{C}_\ell$  is a correspondence defined by

$$\mathcal{C}_\ell(\mathcal{E}^*, D_0) = \{(\bar{Z}, \alpha) : \text{entr}_\ell(\bar{Z}, \alpha, D_0) \geq \mathcal{E}^*, \Lambda_{0j} \leq \mathcal{F}_\ell(\beta_j; \bar{Z}, \alpha) \leq \Upsilon_{0j}\}, \quad (36)$$

<sup>50</sup>The equalities are nonlinear, but that is a minor nuisance. For example, consider the case where  $0 < r < \beta_1$  so that  $k_r = 1$ . Then we must have  $\omega_1 = \omega_2 = \dots = \omega_{J+1}$  and  $\nu_1 = \dots = \nu_{J+1}$ , which leads to the following two equations:  $\alpha\omega_1 \tan\{(v_1 - r)\omega_1\} = -\bar{Z}$  and  $\alpha\omega_1 \tan\{(v_1 - 1)\omega_1\} = 0$ . Our computations have not revealed invertibility problems.

where  $\mathcal{F}_\ell(\beta_j; \bar{Z}, \alpha)$  is the distribution function that has the parametric form described in theorem 9 after partialing out  $\nu, \omega, \tilde{\beta}$  and  $\text{entr}_\ell$  is the entropy level corresponding to  $\mathcal{F}_\ell(\cdot; \bar{Z}, \alpha)$ . So  $\mathcal{F}_\ell(\beta_j; \bar{Z}, \alpha) = \mathcal{F}\{\beta_j; \alpha, \nu_\ell(\bar{Z}, \alpha), \omega_\ell(\bar{Z}, \alpha), \tilde{\beta}_\ell(\bar{Z}, \alpha)\}$ , where  $\nu_\ell, \omega_\ell$ , and  $\tilde{\beta}_\ell$  are defined by recursively solving for  $\nu, \omega$ , and  $\tilde{\beta}$  for given values of  $\bar{Z}, \alpha$ , and the identity  $\ell$  of the bound constraints that hold with equality as explained in appendix E. Note that  $\mathcal{F}_\ell(\beta_j; \cdot, \cdot)$  is a continuous function for all  $j$ . The function  $\text{entr}_\ell(\cdot, \cdot, \cdot)$  can similarly be shown to be continuous.

**Lemma F.1.** For any  $\mathcal{E}^* \in [\underline{\mathcal{E}}^*, \bar{\mathcal{E}}^*]$  and  $D_0, \mathcal{C}_\ell(\mathcal{E}^*, D_0)$  is compact.

*Proof.* Since  $\bar{Z}$  is a probability and  $\alpha$  is a Lagrange multiplier,  $\mathcal{C}_\ell(\mathcal{E}^*, D_0)$  is a bounded set. The closedness of  $\mathcal{C}_\ell(\mathcal{E}^*, D_0)$  follows from continuity of  $\text{entr}_\ell(\cdot, \cdot, \cdot)$  and  $\mathcal{F}_\ell(\beta_j; \cdot, \cdot)$ .  $\square$

**Lemma F.2.** The correspondence  $\mathcal{C}_\ell$  is continuous on  $[\underline{\mathcal{E}}^*, \bar{\mathcal{E}}^*] \times \mathcal{D}$ .

*Proof.* Since  $\text{entr}_\ell(\cdot, \cdot, \cdot)$  and  $\mathcal{F}_\ell(\beta_j; \cdot, \cdot)$  are continuous, by lemma F.1 the graph of  $\mathcal{C}_\ell$  is closed and the image of  $\mathcal{C}_\ell$  is compact. Hence,  $\mathcal{C}_\ell$  is upper-hemicontinuous. We now show that  $\mathcal{C}_\ell$  is also lower-hemicontinuous. Choose an arbitrary point  $(\mathcal{E}^*, D_0, \bar{Z}, \alpha)$  from the graph of  $\mathcal{C}_\ell$ . Let  $(\mathcal{E}_n^*, D_n)$  be an arbitrary sequence in  $[\underline{\mathcal{E}}^*, \bar{\mathcal{E}}^*] \times \mathcal{D}$  that converges to  $(\mathcal{E}^*, D_0)$  as  $n \rightarrow \infty$ . We need to show that there exists  $\{(\bar{Z}_n, \alpha_n) : n = 1, 2, \dots\}$  such that  $(\bar{Z}_n, \alpha_n) \in \mathcal{C}_\ell(\mathcal{E}_n^*, D_n)$  for all sufficiently large  $n$  and for which  $(\bar{Z}_n, \alpha_n) \rightarrow (\bar{Z}, \alpha)$ . Let  $\{\epsilon_n\}$  be a sequence such that  $\epsilon_n \downarrow 0$  and  $\epsilon_n \leq \mathcal{E}^* - \mathcal{E}_n^* \leq 2\epsilon_n$ . Since  $\text{entr}_\ell(\cdot, \cdot, \cdot)$  is continuous at  $(\bar{Z}, \alpha, D_0)$ , there exists  $\delta_n \downarrow 0$  such that

$$(\bar{z}, a, d) \in \mathcal{N}_{\delta_n}(\bar{Z}, \alpha, D_0) \implies |\text{entr}_\ell(\bar{z}, a; d) - \text{entr}_\ell(\bar{Z}, \alpha; D_0)| \leq \epsilon_n, \quad (37)$$

where  $\mathcal{N}_{\delta_n}$  is a  $\delta_n$ -neighborhood. Choose  $n$  large enough to ensure that  $D_n$  is in a  $\delta_n$  neighborhood of  $D_0$ . Then choose  $(\bar{Z}_n, \alpha_n)$  such that  $(\bar{Z}_n, \alpha_n, D_n) \in \mathcal{N}_{\delta_n}(\bar{Z}, \alpha, D_0)$ . It follows that for all such  $n$ ,  $\mathcal{E}^* - \epsilon_n \leq \text{entr}_\ell(\bar{Z}, \alpha, D_0) - \epsilon_n \leq \text{entr}_\ell(\bar{Z}_n, \alpha_n; D_n)$ , where the first inequality holds because  $(\mathcal{E}^*, D_0, \bar{Z}, \alpha)$  belongs to the graph of  $\mathcal{C}_\ell$  and the second inequality follows from (37). Now choose  $\mathcal{E}_n^*$  such that  $\mathcal{E}^* - 2\epsilon_n \leq \mathcal{E}_n^* \leq \mathcal{E}^* - \epsilon_n$ . Then by definition we have  $\mathcal{E}_n^* \leq \mathcal{E}^* - \epsilon_n \leq \text{entr}_\ell(\bar{Z}_n, \alpha_n; D_n)$  and  $(\bar{Z}_n, \alpha_n) \rightarrow (\bar{Z}, \alpha)$ . Repeat the argument for  $\mathcal{F}_\ell$  in lieu of  $\text{entr}_\ell$ .  $\square$

**Proof of theorem 10:** For continuity of  $\pi^\bullet$  it suffices to show that

$$\pi_\ell^\bullet(\mathcal{E}^*, D_0; r) = \min_{(\bar{Z}, \alpha) \in \mathcal{C}(\mathcal{E}^*, D_0)} \pi_\ell^\circ(\bar{Z}, \alpha, \mathcal{E}^*, D_0; r)$$

is a continuous function for each  $\ell$ . Continuity in  $r$  is trivial. Continuity in  $\mathcal{E}^*$  and  $D_0$  follows from the theorem of the maximum because of lemmas F.1 and F.2. Finally, since  $\pi^\bullet$  is continuous, upper hemi-continuity of  $\tilde{r}$  follows from the theorem of the maximum as well.  $\square$

## G. Constraint selection and inference

**Proof of lemma 6.1:** If  $D_0 \in \text{int}(S_{K_0})$ , then by the definition of  $S_{K_0}$  and the KKT conditions,

$$\left\{ \begin{array}{l} \gamma_j^\circ(D_0) > 0, \sum_{k=1}^j g_k^\circ(D_0) = \Upsilon_{0j} \quad \text{for } j \in K_{u0}, \\ \gamma_j^\circ(D_0) = 0, \sum_{k=1}^j g_k^\circ(D_0) < \Upsilon_{0j} \quad \text{for } j \notin K_{u0}, \\ \gamma_j^\circ(D_0) < 0, \sum_{k=1}^j g_k^\circ(D_0) = \Lambda_{j0} \quad \text{for } j \in K_{\ell 0}, \\ \gamma_j^\circ(D_0) = 0, \sum_{k=1}^j g_k^\circ(D_0) > \Lambda_{j0} \quad \text{for } j \notin K_{\ell 0}. \end{array} \right. \quad (38)$$

Therefore,  $K_{u0}^* = K_{u0}$  and  $K_{\ell 0}^* = K_{\ell 0}$  by definition. Now, instead suppose that  $D_0 \in \text{bdr}(S_{K_0})$ . Then, there exist  $j \notin K_{u0} \cup K_{\ell 0}$  and  $\{D_t\}$  with  $D_t \rightarrow D_0$  such that  $\gamma_j^\circ(D_t) \neq 0$  for all  $t$  but  $\gamma_j^\circ(D_t) \rightarrow \gamma_j^\circ(D_0) = 0$ . Fix such  $j$  and  $\{D_t\}$ . By the KKT conditions, for all  $t$  we have either  $\sum_{k=1}^j g_k^\circ(D_t) = \Upsilon_{jt}$  or  $\sum_{k=1}^j g_k^\circ(D_t) = \Lambda_{jt}$ . Therefore, it follows from lemma C.2 that we have either  $\sum_{k=1}^j g_k^\circ(D_0) = \Upsilon_{0j}$  or  $\sum_{k=1}^j g_k^\circ(D_0) = \Lambda_{j0}$ , so  $j \in K_{u0}^* \cup K_{\ell 0}^* = K_0^*$  and hence  $K_0 \neq K_0^*$ .  $\square$

**Proof of lemma 6.2:** Let  $\hat{\gamma}_j^* = \gamma_j^\circ(\hat{D})$  and  $\gamma_{j0}^* = \gamma_j^\circ(D_0)$ . First parts (a) and (b). By theorem 3 and assumption A,  $\hat{Z} = O_p(1)$ . By lemma D.3 and theorem 3, we also have  $\sqrt{N}(\hat{\gamma}_j^* - \gamma_{j0}^*) = O_p(1)$ . Since parts (a) and (b) are similar, we focus on part (a), for which it suffices to show that  $\mathbb{P}(\tilde{K}_u \neq K_{u0}) = o(1)$  and  $\mathbb{P}(\tilde{K}_\ell \neq K_{\ell 0}) = o(1)$ . By symmetry, it suffices to establish the former, where  $\gamma_{j0}^*$  is restricted to be nonnegative. If  $j \notin K_{u0}$  then  $\sqrt{N}\hat{\gamma}_j^* = O_p(1)$ . But then,  $\mathbb{P}(j \in \tilde{K}_u) = \mathbb{P}(\sqrt{N}\hat{\gamma}_j^* > \sqrt{N}\kappa_N) = o(1)$ . Conversely, if  $j \in K_{u0}$  then  $\gamma_{j0}^* > 0$  and  $\sqrt{N}(\hat{\gamma}_j^* - \gamma_{j0}^*) = O_p(1)$ , in which case  $\mathbb{P}(j \notin \tilde{K}_u) = \mathbb{P}(\hat{\gamma}_j^* \leq \kappa_N) = \mathbb{P}\{\sqrt{N}(\hat{\gamma}_j^* - \gamma_{j0}^*) \leq \sqrt{N}(\kappa_N - \gamma_{j0}^*)\} = o(1)$ .

Now the uniform case. For part (a') we note that  $\mathbb{P}(\tilde{K}_u \subset K_{u0}) = 1 - \mathbb{P}(\exists j : j \notin K_{u0}, j \in \tilde{K}_u)$ . But note that  $j \notin K_{u0}$  requires that  $\gamma_{j0}^* = 0$ ; it is insufficient that it converges to 0. Part (b') is similar. For the final assertion, we focus on the upper bounds, and we note that  $\hat{\gamma}_j^* > \kappa_N \Rightarrow \hat{\gamma}_j^* > 0 \Rightarrow \sum_{k=1}^j \hat{g}_k^* = \hat{\Upsilon}_j \Rightarrow \sum_{k=1}^j \hat{g}_k^* > \hat{\Upsilon}_j - \kappa_N$ . The argument for the lower bounds is similar.  $\square$

**Proof of theorem 4:** By lemma 6.1,  $\mathbb{P}(\hat{T} \neq T) = o(1)$ . Suppose first that  $D_0$  is a boundary point, such that by lemma 6.1  $K_0 \subsetneq K_0^*$ . By theorem 3 and lemma 6.2, and the continuity of  $G$  in its first

argument,

$$\begin{aligned}\mathbb{P}(\hat{Z} \leq x) &= \mathbb{P}\{\Theta(\hat{\Omega}, \hat{K}) > x\} + o(1) \geq \mathbb{P}\left\{\max_{\tilde{K} \subseteq K \subseteq \tilde{K}^*} \Theta(\hat{\Omega}, K) \leq x\right\} + o(1) \\ &= \mathbb{P}\left\{\max_{K_0 \subseteq K \subseteq K_0^*} \Theta(\hat{\Omega}, K) \leq x\right\} + o(1) = \mathbb{P}(T \leq x) + o(1).\end{aligned}$$

Now, suppose instead that  $D_0$  is an interior point. Lemma 6.1 implies that then  $K_0 = K_0^*$ , such that  $\mathbb{P}(\hat{Z} \leq x) = \mathbb{P}\{\Theta(\Phi, K_0) \leq x\} + o(1) = \mathbb{P}(T \leq x) + o(1)$ .  $\square$

## H. Optimal reserve price and maximum revenue

### H.1 Consistency of $\hat{\mathcal{R}}_N$ :

**Proof of lemma 6.3:** Follows from the fact that  $\pi(\cdot, g^*)$  is continuous on  $[0, 1]$  and (strictly) concave on each  $I_j$ .  $\square$

**Lemma H.1.**  $\sup_{v \in [0, 1]} |\hat{\ell}^*(v) - \ell^*(v)| + \sup_{v \in [0, 1]} |\hat{\mathcal{F}}^*(v) - \mathcal{F}^*(v)| = O_p(1/\sqrt{N})$ .

*Proof.* It follows from (5) and (6) and theorem 3.  $\square$

**Lemma H.2.**  $\sup_{r \in [0, 1]} |\pi(r, \hat{g}^*) - \pi(r, g^*)| = O_p(1/\sqrt{N})$ .

*Proof.* It follows from lemma H.1 and (1).  $\square$

**Lemma H.3.**  $|\mathcal{P}(\hat{g}^*) - \mathcal{P}(g^*)| = O_p(1/\sqrt{N})$

*Proof.* Follows from lemma H.2.  $\square$

**Proof of theorem 5:** Note that by continuity of  $\pi(\cdot, g^*)$  and by lemma 6.3,  $\mathcal{R}_\epsilon = \{r : \pi(r, g^*) \geq \mathcal{P}(g^*) - \epsilon\}$  consists for any sufficiently small  $\epsilon > 0$  of a union of disjoint compact intervals  $\mathcal{R}_{\epsilon 1}, \dots, \mathcal{R}_{\epsilon m}$  each containing one  $r_j^*$ . Choose  $\epsilon > 0$ . By lemmas H.2 and H.3,

$$\begin{cases} \mathbb{P}(\hat{\mathcal{R}}_N \not\subseteq \mathcal{R}_\epsilon) \leq \mathbb{P}\left[\max_{r \in [0, 1]} \{\pi(r, \hat{g}^*) - \pi(r, g^*)\} \geq \mathcal{P}(\hat{g}^*) - \mathcal{P}(g^*) + \epsilon - \kappa_n\right] = o(1), \\ \mathbb{P}(\mathcal{R}_0 \not\subseteq \hat{\mathcal{R}}_N) \leq \mathbb{P}\left[\max_{r \in [0, 1]} \{\pi(r, g^*) - \pi(r, \hat{g}^*)\} \geq \mathcal{P}(g^*) - \mathcal{P}(\hat{g}^*) + \kappa_n\right] = o(1).\end{cases}$$

Let  $\epsilon \rightarrow 0$ .  $\square$

## H.2 Sensitivity

**Proof of lemma 6.4:** Note that  $r_1(\hat{g}^*) \neq \hat{r}_1 \Rightarrow r_1(\hat{g}^*) \notin \hat{\mathcal{R}}_{N,1}$ . Now,  $r_1(\hat{g}^*) \in \hat{\mathcal{R}}_N$  with probability approaching one because otherwise  $\mathcal{P}(\hat{g}^*) - \kappa_N \leq \pi(\hat{r}_1, \hat{g}^*) \leq \pi\{r_1(\hat{g}^*), \hat{g}^*\} < \mathcal{P}(\hat{g}^*) - \kappa_N$ , which is a contradiction. But since  $r_1(\hat{g}^*) \xrightarrow{p} r_1^*$  by construction,  $r_1(\hat{g}^*) \in \hat{\mathcal{R}}_{N,1}$  because  $\hat{\mathcal{R}}_{N,1}$  is the collection of elements in  $\hat{\mathcal{R}}_N$  that converges to  $r_1^*$ .  $\square$

**Lemma H.4.** If  $\hat{r}_* - r_k^* = o_p(1)$  for some  $\hat{r}_* \in \hat{\mathcal{R}}_N$  and some  $k = 1, \dots, m$  then  $\hat{r}_* - r_k^* = O_p(\sqrt{\kappa_N})$ .

*Proof.* Let  $j^*$  be such that  $r_k^* \in I_{j^*}$ . If  $r_k^* > \beta_{j^*-1}$  then  $O_p(\kappa_N) = \pi(\hat{r}_*, g^*) - \pi(r_k^*, g^*) \simeq \partial_r^2 \pi(r_k^*, g^*)(\hat{r}_* - r_k^*)^2 / 2$ .<sup>51</sup> Since  $\partial_r^2 \pi(r_k^*, g^*) < 0$  it follows that  $\hat{r}_* - r_k^* = O_p(\sqrt{\kappa_N})$ . If  $r_k^* = \beta_{j^*-1}$  then one should consider  $\partial_r^+ \pi(r_k^*, g^*)$  and  $\partial_r^- \pi(r_k^*, g^*)$  separately. If both are nonzero then it follows that  $\hat{r}_* - r_k^* = O_p(\kappa_N)$ . If either the right or left derivative equals zero then the convergence rate is no worse than  $\sqrt{\kappa_N}$ , as before.  $\square$

**Proof of theorem 6:** Suppose first that  $m = 1$ . Consistency of  $\hat{r}_m$  follows trivially from theorem 5 and lemma H.4 establishes that  $\hat{r}_m - r_m^* = O_p(\sqrt{\kappa_N})$ . We now argue that  $\hat{m} = m$  with probability approaching one. Suppose that  $\hat{m} > m$ . Then  $\hat{r}_{m+1} \xrightarrow{p} r_m^*$  by theorem 5 and hence  $\hat{r}_{m+1} - r_m^* = O_p(\sqrt{\kappa_N})$  by lemma H.4. But the construction of  $\hat{\mathcal{R}}_{N,m+1}$  implies that

$$\sqrt[3]{\kappa_N} = \tilde{\kappa}_N \leq \hat{r}_{m+1} - \min \hat{\mathcal{R}}_{N,m} = \hat{r}_{m+1} - r_m^* + r_m^* - \min \hat{\mathcal{R}}_{N,m} \leq \hat{r}_{m+1} - r_m^* + O_p(\sqrt{\kappa_N}),$$

which is at odds with  $\hat{r}_{m+1} - r_m^* = O_p(\sqrt{\kappa_N})$ . So  $\hat{m} = m$  with probability approaching one and the proof is complete for  $m = 1$ .

Now suppose that  $m = 2$ . Consistency and convergence rate of  $\hat{r}_{m-1}$  follow as above. Further,  $\hat{m} \geq m$  with probability approaching one since  $\max \hat{\mathcal{R}}_N \xrightarrow{p} r_m^*$ . As established above  $\hat{r}_m$  does not converge to  $r_{m-1}^*$ , so  $\hat{r}_m$  must converge to  $r_m^*$  and hence  $\hat{r}_m - r_m^* = O_p(\sqrt{\kappa_N})$ . Using the same argument as when  $m = 1$ , now  $\hat{m} = m$  with probability approaching one. Iterate the argument made for  $m = 2$  for  $m > 2$ , noting that  $m$  is finite.  $\square$

**H.2 Sensitivity:** Fixing a direction  $d$ , let  $\Psi(d, t) = [r_1\{g^\circ(D_0 + td)\} - r_1\{g^\circ(D_0)\}]/t$ . The existence of  $\lim_{t \downarrow 0} \Psi(d, t)$  requires the Hadamard differentiability of  $r_1$  at  $g^* = g^\circ(D_0)$ : the chain rule fails if  $r_1$  is only directionally differentiable. So, we first consider the Hadamard directional derivative of  $r_1$ : for any  $v_t = v + o(1)$ ,  $\nabla_H r_1(g^*, v) = \lim_{t \downarrow 0} \{r_1(g^* + tv_t) - r_1(g^*)\}/t$ .

**Lemma H.5.**  $r_1$  is continuous at  $g^*$ .

*Proof.* By the maximum theorem,  $\tilde{R}_1(g) = \operatorname{argmax}_{r \in A_1} \pi(r, g)$  is upper hemicontinuous. So, the conclusion follows from the fact that  $\tilde{R}_1(g^*)$  is a singleton.  $\square$

<sup>51</sup>We define  $\simeq$  to mean that any remaining terms are asymptotically negligible.

## H.2 Sensitivity

**Lemma H.6.** For cases I–IV,  $\nabla_H r_1(g^*, v) = \psi_{g^*}(v)$ . First case I. By lemma H.5, we must have  $\beta_{j^*-1} < r_1(g) < \beta_{j^*}$  in a neighborhood of  $g^*$ . Hence  $\theta\{r_1(g), g\} = 0$  in a neighborhood of  $g^*$ . Apply the implicit function theorem.

For case II, note that  $\theta(\cdot, g)$  is an invertible function near  $r_1^* = \beta_{j^*-1}$ , that both  $\theta$  and its inverse are continuous in  $g$ , and that  $\theta$ 's inverse is flat in its first argument. For cases III and IV, combine the arguments for cases I and II.  $\square$

**Lemma H.7.** For case V and any  $v_t = v + o(1)$ , we have  $\limsup_{t \downarrow 0} |r_1(g^* + tv_t) - r_1(g^*)| / t \leq \max(|v^\top \delta|, |v^\top \delta^-|)$ .

*Proof.* Follows immediately by noting that

$$\limsup_{t \downarrow 0} \frac{|r_1(g^* + tv_t) - r_1(g^*)|}{t} \leq \max \left\{ \left| \frac{\partial_g \theta^+(r^*, g^*)}{\partial_r \theta^+(r^*, g^*)} \right|, \left| \frac{\partial_g \theta^-(r^*, g^*)}{\partial_r \theta^-(r^*, g^*)} \right| \right\}. \quad \square$$

**Lemma H.8.** In cases I–IV we have  $\nabla(r_1 \circ g^*)(D_0, d) = \nabla_H r_1\{g^*, \nabla g^\circ(D_0, d)\}$ , where  $\nabla_H r_1(g^*, v)$  is given in lemma H.6. For V, we have  $\limsup_{t \downarrow 0} |\Psi(d, t)| \leq \max\{|\nabla g^\circ(D_0, d)^\top \delta|, |\nabla g^\circ(D_0, d)^\top \delta^-|\}$ .

*Proof.* Follows from lemmas D.2 and H.6.  $\square$

$$\text{Let } \tilde{d} = (\hat{D} - D_0) / \|\hat{D} - D_0\|.$$

**Lemma H.9.** In cases I–IV, we have

$$\sqrt{N} \{r_1(\hat{g}^*) - r_1(g^*)\} = \nabla_H r_1\{g^*, \nabla g^\circ(D_0, \tilde{d})\} \sqrt{N} \|\hat{D} - D_0\| + o_p(1),$$

whereas in case V,

$$\sqrt{N} |r_1(\hat{g}^*) - r_1(g^*)| \leq \max\{|\nabla g^\circ(D_0, \tilde{d})^\top \delta|, |\nabla g^\circ(D_0, \tilde{d})^\top \delta^-|\} \sqrt{N} \|\hat{D} - D_0\| + o_p(1).$$

*Proof.* Note that  $r_1(\hat{g}^*) = r_1\{g^* + \tilde{d} \|\hat{D} - D_0\|\}$ . Using lemma H.8, apply the same logic as the proof of theorem 3.  $\square$

**Proof of theorem 7:** Follows from lemmas 6.4 and H.9.  $\square$

**Proof of theorem 8:** The function  $\pi$  is differentiable in  $g$  and directionally differentiable in  $r$ , so

$$\begin{aligned} \pi(r, g) - \pi(r_1^*, g^*) &= \partial_r^+ \pi(r_1^*, g^*) \max(0, r - r_1^*) + \partial_r^- \pi(r_1^*, g^*) \min(0, r - r_1^*) \\ &\quad + \partial_{g^\top} \pi(r_1^*, g^*) (g - g^*) + o(|r - r_1^*| + \|g - g^*\|), \end{aligned}$$

## REFERENCES

where  $\partial_r^+$  and  $\partial_r^-$  denote the right and left derivative, respectively. Therefore, by theorems 3 and 7,

$$\begin{aligned} \sqrt{N}\{\pi(\hat{r}_1, \hat{g}^*) - \pi(r_1^*, g^*)\} &= \partial_r^+ \pi(r_1^*, g^*) \max\{0, \sqrt{N}(\hat{r}_1 - r_1^*)\} \\ &+ \partial_r^- \pi(r_1^*, g^*) \min\{0, \sqrt{N}(\hat{r}_1 - r_1^*)\} + \partial_{g^\top} \pi(r_1^*, g^*) \sqrt{N}(\hat{g}^* - g^*) + o_p(1). \end{aligned} \quad (39)$$

Now, note that by theorem 7,

1.  $\partial_r^+ \pi(r_1^*, g^*) = 0 = \partial_r^- \pi(r_1^*, g^*)$  in cases I and V;
2.  $\sqrt{N}(\hat{r}_1 - r_1^*) = o_p(1)$  in case II;
3.  $\partial_r^+ \pi(r_1^*, g^*) = 0$  and  $\sqrt{N}(\hat{r}_1 - r_1^*) - o_p(1) \geq 0$  in case III;
4.  $\partial_r^- \pi(r_1^*, g^*) = 0$  and  $\sqrt{N}(\hat{r}_1 - r_1^*) - o_p(1) \leq 0$  in case IV.

Therefore, the first right-hand side terms in (39) are asymptotically negligible in all cases.  $\square$

## References

- Andrews, Donald WK and Gustavo Soares (2010). “Inference for parameters defined by moment inequalities using generalized moment selection”. In: *Econometrica* 78.1, pp. 119–157.
- Aryal, Gaurab and Dong-Hyuk Kim (2013). “A point decision for partially identified auction models”. In: *Journal of Business & Economic Statistics* 31.4, pp. 384–397.
- Athey, Susan and Philip A Haile (2002). “Identification of standard auction models”. In: *Econometrica* 70.6, pp. 2107–2140.
- Avery, Christopher (1998). “Strategic jump bidding in English auctions”. In: *The Review of Economic Studies* 65.2, pp. 185–210.
- Bertsekas, Dimitri P (2015). *Convex optimization algorithms*. Athena Scientific Belmont.
- Bulow, Jeremy and Paul Klemperer (1996). “Auctions versus negotiations”. In: *The American Economic Review*, pp. 180–194.
- Chernozhukov, Victor, Han Hong, and Elie Tamer (2007). “Estimation and confidence regions for parameter sets in econometric models”. In: *Econometrica* 75.5, pp. 1243–1284.
- Chesher, Andrew and Adam Rosen (2018). *Econometric analysis of incomplete English auction models*. Tech. rep.
- Coey, Dominic et al. (2017). “Ascending auctions with bidder asymmetries”. In: *Quantitative Economics* 8.1, pp. 181–200.
- Daniel, Kent D and David A Hirshleifer (1998). *A theory of costly sequential bidding*. Tech. rep.
- Gilboa, Itzhak and David Schmeidler (1989). “Maxmin expected utility with non-unique prior”. In: *Journal of mathematical economics* 18.2, pp. 141–153.

## REFERENCES

- Golan, Amos (2018). *Foundations of info-metrics: Modeling, inference, and imperfect information*. Oxford University Press.
- Grünwald, Peter D and A Philip Dawid (2004). “Game theory, maximum entropy, minimum discrepancy and robust Bayesian decision theory”. In: *Annals of Statistics*, pp. 1367–1433.
- Haile, Philip A and Elie Tamer (2003). “Inference with an incomplete model of English auctions”. In: *Journal of Political Economy* 111.1, pp. 1–51.
- Hansen, Lars Peter and Thomas J Sargent (2008). *Robustness*. Princeton university press.
- Hardle, Wolfgang, Peter Hall, and Hidehiko Ichimura (1993). “Optimal smoothing in single-index models”. In: *The annals of Statistics*, pp. 157–178.
- Harremoës, Peter and Flemming Topsøe (2001). “Maximum entropy fundamentals”. In: *Entropy* 3.3, pp. 191–226.
- Hartline, Jason D (2016). *Mechanism design and approximation*. draft.
- Hsieh, Yu-Wei, Xiaoxia Shi, and Matthew Shum (2018). “Inference on estimators defined by mathematical programming”. In: *Available at SSRN 3041040*.
- Ichimura, Hidehiko (1993). “Semiparametric least squares (SLS) and weighted SLS estimation of single-index models”. In: *Journal of Econometrics* 58.1-2, pp. 71–120.
- Jun, Sung Jae and Joris Pinkse (2016). *Point prediction under partial identification*. Tech. rep. Penn State.
- Larsen, Bradley J (2014). “The efficiency of real-world bargaining: Evidence from wholesale used-auto auctions”. In: *The Review of Economic Studies*.
- Myerson, Roger B (1981). “Optimal auction design”. In: *Mathematics of operations research* 6.1, pp. 58–73.
- Topsøe, Flemming (1979). “Information-theoretical optimization techniques”. In: *Kybernetika* 15.1, pp. 8–27.